

WO 2004/041170

PCT/US2003/034312

COMPOSITIONS AND METHODS FOR THE TREATMENT OF IMMUNE RELATED DISEASES

5

Field of the Invention

The present invention relates to compositions and methods useful for the diagnosis and treatment of immune related diseases.

Background of the Invention

10

Immune related and inflammatory diseases are the manifestation or consequence of fairly complex, often multiple interconnected biological pathways which in normal physiology are critical to respond to insult or injury, initiate repair from insult or injury, and mount innate and acquired defense against foreign organisms. Disease or pathology occurs when these normal physiological pathways cause additional insult or injury either as directly related to the intensity of the response, as a consequence of abnormal regulation or excessive stimulation, as a reaction to self, or as a combination of these.

15

Though the genesis of these diseases often involves multistep pathways and often multiple different biological systems/pathways, intervention at critical points in one or more of these pathways can have an ameliorative or therapeutic effect. Therapeutic intervention can occur by either antagonism of a detrimental process/pathway or stimulation of a beneficial process/pathway.

20

Many immune related diseases are known and have been extensively studied. Such diseases include immune-mediated inflammatory diseases, non-immune-mediated inflammatory diseases, infectious diseases, immunodeficiency diseases, neoplasia, etc.

Immune related diseases could be treated by suppressing the immune response. Using neutralizing antibodies that inhibit molecules having immune stimulatory activity would be beneficial in the treatment of immune-mediated and inflammatory diseases. Molecules which inhibit the immune response can be utilized (proteins directly or via the use of antibody agonists) to inhibit the immune response and thus ameliorate immune related disease.

25

Macrophages represent an ubiquitously distributed population of fixed and circulating mononuclear phagocytes that express a variety of functions including cytokine production, killing of microbes and tumor cells and processing and presentation of antigens. Macrophages originate in the bone marrow from stem cells that give rise to a bipotent granulocyte/macrophage cell population. Distinct granulocyte and macrophage colony forming cell lineages arise from GM-CSF under the influence of specific cytokines. Upon division, monoblasts give rise to promonocytes and monocytes in the bone marrow. From there, monocytes enter the circulation. In response to particular stimuli (e.g. infection or foreign bodies) monocytes migrate into tissues and organs where they differentiate into macrophages.

30

35

Macrophages in various tissues vary in their morphology and function and have been assigned different names, e.g. Kupffer cells in the liver, pulmonary and alveolar macrophages in the lung and microglial cells in the central nervous system. However, the relationship between blood monocytes and tissue macrophages remains unclear.

In the present study, monocytes were differentiated into macrophages by adherence to plastic in the presence of a combination of human and bovine serum. After 7 days in culture, monocytes-derived macrophages display features typical of differentiated tissue macrophages including their ability to phagocytose opsonized particles, secretion of TNF-alpha upon lipopolysaccharide (LPS) stimulation, formation of processes and the presence of macrophage cell surface markers.

Using microarray technologies, gene transcripts from non-differentiated monocytes harvested before adhering were compared with those at 1 day and 7 days in culture. Genes selectively expressed in monocytes or macrophages could be used for the diagnosis and treatment of various chronic inflammatory or autoimmune diseases in the human. In particular, surface expressed molecules or transmembrane receptors involved in monocyte/macrophage adhesion and endothelial cell transmigration could provide novel targets to treat chronic inflammation by interference with the homing of these cells to the site of inflammation. In addition, transmembrane inhibitory receptors could be used to down-regulate monocyte/macrophage effector functions. Therapeutic molecules can be antibodies, peptides, fusion proteins or small molecules.

Despite the above research in monocyte/macrophages, there is a great need for additional diagnostic and therapeutic agents capable of detecting the presence of monocyte/macrophage mediated disorders in a mammal and for effectively reducing these disorders. Accordingly, it is an objective of the present invention to identify polypeptides that are differentially expressed in macrophages as compared to non-differentiated monocytes, and to use those polypeptides, and their encoding nucleic acids, to produce compositions of matter useful in the therapeutic treatment and diagnostic detection of monocyte/macrophage mediated disorders in mammals.

#### Summary of the Invention

##### A. Embodiments

The present invention concerns compositions and methods useful for the diagnosis and treatment of immune related disease in mammals, including humans. The present invention is based on the identification of proteins (including agonist and antagonist antibodies) which are a result of stimulation of the immune response in mammals. Immune related diseases can be treated by suppressing or enhancing the immune response. Molecules that enhance the immune response stimulate or potentiate the immune response to an antigen. Molecules which stimulate the immune response can be used therapeutically where enhancement of the immune response would be beneficial. Alternatively, molecules that suppress the immune response attenuate or reduce the immune response to an antigen (e.g., neutralizing antibodies) can be used therapeutically where attenuation of the immune response would be beneficial (e.g., inflammation). Accordingly, the PRO polypeptides, agonists and antagonists thereof are also useful to prepare medicines and medicaments for the treatment of immune-related and inflammatory diseases. In a specific aspect, such medicines and medicaments comprise a therapeutically effective amount of a PRO polypeptide, agonist or antagonist thereof with a pharmaceutically acceptable carrier. Preferably, the admixture is sterile.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprises contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a

native-sequence-PRO polypeptide. In a specific aspect, the PRO agonist or antagonist is an anti-PRO antibody.

In another embodiment, the invention concerns a composition of matter comprising a PRO polypeptide or an agonist or antagonist antibody which binds the polypeptide in admixture with a carrier or excipient. In one aspect, the composition comprises a therapeutically effective amount of the polypeptide or antibody. In another aspect, when the composition comprises an immune stimulating molecule, the composition is useful for: (a) increasing infiltration of inflammatory cells into a tissue of a mammal in need thereof, (b) stimulating or enhancing an immune response in a mammal in need thereof, (c) increasing the proliferation of monocytes/macrophages in a mammal in need thereof in response to an antigen, (d) stimulating the activity of monocytes/macrophages or (e) increasing the vascular permeability. In a further aspect, when the composition comprises an immune inhibiting molecule, the composition is useful for: (a) decreasing infiltration of inflammatory cells into a tissue of a mammal in need thereof, (b) inhibiting or reducing an immune response in a mammal in need thereof, (c) decreasing the activity of monocytes/macrophages or (d) decreasing the proliferation of monocytes/macrophages in a mammal in need thereof in response to an antigen. In another aspect, the composition comprises a further active ingredient, which may, for example, be a further antibody or a cytotoxic or chemotherapeutic agent. Preferably, the composition is sterile.

In another embodiment, the invention concerns a method of treating an immune related disorder in a mammal in need thereof, comprising administering to the mammal an effective amount of a PRO polypeptide, an agonist thereof, or an antagonist thereto. In a preferred aspect, the immune related disorder is selected from the group consisting of: systemic lupus erythematosus, rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, spondyloarthropathies, systemic sclerosis, idiopathic inflammatory myopathies, Sjögren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia, autoimmune thrombocytopenia, thyroiditis, diabetes mellitus, immune-mediated renal disease, demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious, autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory bowel disease, gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft -versus-host-disease.

In another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody. In one aspect, the present invention concerns an isolated antibody which binds a PRO polypeptide. In another aspect, the antibody mimics the activity of a PRO polypeptide (an agonist antibody) or conversely the antibody inhibits or neutralizes the activity of a PRO polypeptide (an antagonist antibody). In another aspect, the antibody is a monoclonal antibody, which preferably has nonhuman complementarity determining region (CDR) residues and human framework region

(FR) residues. The antibody may be labeled and may be immobilized on a solid support. In a further aspect, the antibody is an antibody fragment, a monoclonal antibody, a single-chain antibody, or an anti-idiotypic antibody.

In yet another embodiment, the present invention provides a composition comprising an anti-PRO antibody in admixture with a pharmaceutically acceptable carrier. In one aspect, the composition comprises a therapeutically effective amount of the antibody. Preferably, the composition is sterile. The composition may be administered in the form of a liquid pharmaceutical formulation, which may be preserved to achieve extended storage stability. Alternatively, the antibody is a monoclonal antibody, an antibody fragment, a humanized antibody, or a single-chain antibody.

10 In a further embodiment, the invention concerns an article of manufacture, comprising:

- (a) a composition of matter comprising a PRO polypeptide or agonist or antagonist thereof;
- (b) a container containing said composition; and
- (c) a label affixed to said container, or a package insert included in said container referring to the use of said PRO polypeptide or agonist or antagonist thereof in the treatment of an immune related

15 disease. The composition may comprise a therapeutically effective amount of the PRO polypeptide or the agonist or antagonist thereof.

In yet another embodiment, the present invention concerns a method of diagnosing an immune related disease in a mammal, comprising detecting the level of expression of a gene encoding a PRO polypeptide (a) in a test sample of tissue cells obtained from the mammal, and (b) in a control sample of

20 known normal tissue cells of the same cell type, wherein a higher or lower expression level in the test sample as compared to the control sample indicates the presence of immune related disease in the mammal from which the test tissue cells were obtained.

In another embodiment, the present invention concerns a method of diagnosing an immune disease in a mammal, comprising (a) contacting an anti-PRO antibody with a test sample of tissue cells obtained

25 from the mammal, and (b) detecting the formation of a complex between the antibody and a PRO polypeptide, in the test sample; wherein the formation of said complex is indicative of the presence or absence of said disease. The detection may be qualitative or quantitative, and may be performed in comparison with monitoring the complex formation in a control sample of known normal tissue cells of the same cell type. A larger quantity of complexes formed in the test sample indicates the presence or absence

30 of an immune disease in the mammal from which the test tissue cells were obtained. The antibody preferably carries a detectable label. Complex formation can be monitored, for example, by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. The test sample is usually obtained from an individual suspected of having a deficiency or abnormality of the immune system.

In another embodiment, the invention provides a method for determining the presence of a PRO polypeptide in a sample comprising exposing a test sample of cells suspected of containing the PRO polypeptide to an anti-PRO antibody and determining the binding of said antibody to said cell sample. In a

35 specific aspect, the sample comprises a cell suspected of containing the PRO polypeptide and the antibody binds to the cell. The antibody is preferably detectably labeled and/or bound to a solid support.

In another embodiment, the present invention concerns an immune-related disease diagnostic kit,

40 comprising an anti-PRO antibody and a carrier in suitable packaging. The kit preferably contains



instructions for using the antibody to detect the presence of the PRO polypeptide. Preferably the carrier is pharmaceutically acceptable.

5 In another embodiment, the present invention concerns a diagnostic kit, containing an anti-PRO antibody in suitable packaging. The kit preferably contains instructions for using the antibody to detect the PRO polypeptide.

In another embodiment, the invention provides a method of diagnosing an immune-related disease in a mammal which comprises detecting the presence or absence of a PRO polypeptide in a test sample of tissue cells obtained from said mammal, wherein the presence or absence of the PRO polypeptide in said test sample is indicative of the presence of an immune-related disease in said mammal.

10 In another embodiment, the present invention concerns a method for identifying an agonist of a PRO polypeptide comprising:

(a) contacting cells and a test compound to be screened under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and

(b) determining the induction of said cellular response to determine if the test compound is an effective agonist, wherein the induction of said cellular response is indicative of said test compound being an effective agonist.

15 In another embodiment, the invention concerns a method for identifying a compound capable of inhibiting the activity of a PRO polypeptide comprising contacting a candidate compound with a PRO polypeptide under conditions and for a time sufficient to allow these two components to interact and determining whether the activity of the PRO polypeptide is inhibited. In a specific aspect, either the candidate compound or the PRO polypeptide is immobilized on a solid support. In another aspect, the non-immobilized component carries a detectable label. In a preferred aspect, this method comprises the steps of:

(a) contacting cells and a test compound to be screened in the presence of a PRO polypeptide under conditions suitable for the induction of a cellular response normally induced by a PRO polypeptide; and

25 (b) determining the induction of said cellular response to determine if the test compound is an effective antagonist.

In another embodiment, the invention provides a method for identifying a compound that inhibits the expression of a PRO polypeptide in cells that normally express the polypeptide, wherein the method comprises contacting the cells with a test compound and determining whether the expression of the PRO polypeptide is inhibited. In a preferred aspect, this method comprises the steps of:

30 (a) contacting cells and a test compound to be screened under conditions suitable for allowing expression of the PRO polypeptide; and

(b) determining the inhibition of expression of said polypeptide.

35 In yet another embodiment, the present invention concerns a method for treating an immune-related disorder in a mammal that suffers therefrom comprising administering to the mammal a nucleic acid molecule that codes for either (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide or (c) an antagonist of a PRO polypeptide, wherein said agonist or antagonist may be an anti-PRO antibody. In a preferred embodiment, the mammal is human. In another preferred embodiment, the nucleic acid is administered via *ex vivo* gene therapy. In a further preferred embodiment, the nucleic acid is comprised within a vector, more preferably an adenoviral, adeno-associated viral, lentiviral or retroviral vector.

In yet another aspect, the invention provides a recombinant viral particle comprising a viral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide, or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein the viral vector is in association with viral structural proteins. Preferably, the signal sequence is from a mammal, such as from a native PRO polypeptide.

In a still further embodiment, the invention concerns an *ex vivo* producer cell comprising a nucleic acid construct that expresses retroviral structural proteins and also comprises a retroviral vector consisting essentially of a promoter, nucleic acid encoding (a) a PRO polypeptide, (b) an agonist polypeptide of a PRO polypeptide or (c) an antagonist polypeptide of a PRO polypeptide, and a signal sequence for cellular secretion of the polypeptide, wherein said producer cell packages the retroviral vector in association with the structural proteins to produce recombinant retroviral particles.

In a still further embodiment, the invention provides a method of increasing the activity of monocytes/macrophages in a mammal comprising administering to said mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein the activity of monocytes/macrophages in the mammal is increased.

In a still further embodiment, the invention provides a method of decreasing the activity of monocytes/macrophages in a mammal comprising administering to said mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein the activity of monocytes/macrophages in the mammal is decreased.

In a still further embodiment, the invention provides a method of increasing the proliferation of monocytes/macrophages in a mammal comprising administering to said mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein the proliferation of monocytes/macrophages in the mammal is increased.

In a still further embodiment, the invention provides a method of decreasing the proliferation of monocytes/macrophages in a mammal comprising administering to said mammal (a) a PRO polypeptide, (b) an agonist of a PRO polypeptide, or (c) an antagonist of a PRO polypeptide, wherein the proliferation of monocytes/macrophages in the mammal is decreased.

#### B. Additional Embodiments

In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

5 In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

In other embodiments, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

10 In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

25 In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an

extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a  
 5 nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity,  
 10 alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity,  
 15 alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs as disclosed herein, or (b) the complement of the DNA molecule of (a).

Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide  
 20 sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the  
 25 complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60  
 30 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160  
 35 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500  
 40 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700

nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences herein above identified.

In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid

sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs as disclosed herein.

5 In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as herein before described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

10 Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

15 In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

25 Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as herein before described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

#### 30 BRIEF DESCRIPTION OF THE DRAWINGS

In the list of figures for the present application, specific cDNA sequences which are differentially expressed in differentiated macrophages as compared to normal undifferentiated monocytes are individually identified with a specific alphanumerical designation. These cDNA sequences are differentially expressed in monocytes that are specifically treated as described in Example 1 below. If start and/or stop codons have been identified in a cDNA sequence shown in the attached figures, they are shown in bold and underlined font, and the encoded polypeptide is shown in the next consecutive figure.

35 The Figures 1-2517 show the nucleic acids of the invention and their encoded PRO polypeptides. Also included, for convenience is a List of Figures attached hereto as Appendix A, which gives the figure number and the corresponding DNA or PRO number.

## List of Figures

- Figure 1: DNA227321, NP\_001335.1, 200046.at  
 Figure 2: PRO37784  
 Figure 3: DNA304680, HSPCB, 200064.at  
 Figure 4: PRO71106  
 Figure 5: DNA328347, NP\_002146.1, 117.at  
 Figure 6: PRO58142  
 Figure 7A-B: DNA328348, MAP4, 243.g.at  
 Figure 8: PRO84209  
 Figure 9: DNA83128, NP\_002979.1, 32128.at  
 Figure 10: PRO2601  
 Figure 11: DNA272223, NP\_004444.1, 33494.at  
 Figure 12: PRO60485  
 Figure 13: DNA327522, NP\_000396.1, 33646.g.at  
 Figure 14: PRO2874  
 Figure 15: DNA328349, NP\_004556.1, 33760.at  
 Figure 16: PRO84210  
 Figure 17A-B: DNA328350, NP\_056155.1, 34764.at  
 Figure 18: PRO84211  
 Figure 19: DNA328351, NP\_006143.1, 35974.at  
 Figure 20: PRO84212  
 Figure 21: DNA328352, NP\_004183.1, 36553.at  
 Figure 22: PRO84213  
 Figure 23: DNA271996, NP\_004928.1, 36566.at  
 Figure 24: PRO60271  
 Figure 25: DNA326969, NP\_036455.1, 36711.at  
 Figure 26: PRO83282  
 Figure 27: DNA304703, NP\_005923.1, 36830.at  
 Figure 28: PRO71129  
 Figure 29: DNA328353, AAB72234.1, 37079.at  
 Figure 30: PRO84214  
 Figure 31: DNA103289, NP\_006229.1, 37152.at  
 Figure 32: PRO4619  
 Figure 33A-B: DNA255096, NP\_055449.1, 37384.at  
 Figure 34: PRO50180  
 Figure 35: DNA256295, NP\_002310.1, 37796.at  
 Figure 36: PRO51339  
 Figure 37: DNA328354, PARVB, 37965.at  
 Figure 38: PRO84215  
 Figure 39: DNA53531, NP\_001936.1, 38037.at  
 Figure 40: PRO131  
 Figure 41: DNA254127, NP\_008925.1, 38241.at  
 Figure 42: PRO49242  
 Figure 43: DNA328355, NP\_006471.2, 38290.at  
 Figure 44: PRO84216  
 Figure 45: DNA328356, BC013566, 39248.at  
 Figure 46: PRO38028  
 Figure 47: DNA328357, 1452321.2, 39582.at  
 Figure 48: PRO84217  
 Figure 49A-B: DNA328358, STK10, 40420.at  
 Figure 50: PRO84218  
 Figure 51A-B: DNA328359, BAA21572.1, 41386.i.at  
 Figure 52: PRO84219  
 Figure 53A-D: DNA328360, NP\_055061.1, 41660.at  
 Figure 54: PRO84220  
 Figure 55: DNA327526, BC001698, 45288.at  
 Figure 56: PRO83574  
 Figure 57A-B: DNA328361, BAA92570.1, 47773.at  
 Figure 58: PRO84221  
 Figure 59: DNA328362, NP\_060312.1, 48106.at  
 Figure 60: PRO84222  
 Figure 61: DNA328363, DNA328363, 52651.at  
 Figure 62: PRO84685  
 Figure 63: DNA328364, NP\_068577.1, 52940.at  
 Figure 64: PRO84223  
 Figure 65A-B: DNA327528, BAB33338.1, 55081.at  
 Figure 66: PRO83576  
 Figure 67: DNA225650, NP\_057246.1, 48825.at  
 Figure 68: PRO36113  
 Figure 69: DNA328365, NP\_060541.1, 58780.s.at  
 Figure 70: PRO84224  
 Figure 71: DNA328366, NP\_079233.1, 59375.at  
 Figure 72: PRO84225  
 Figure 73: DNA328367, NP\_079108.2, 60471.at  
 Figure 74: PRO84226  
 Figure 75: DNA327876, NP\_005081.1, 60528.at  
 Figure 76: PRO83815  
 Figure 77A-B: DNA328368, 1503444.3, 87100.at  
 Figure 78: PRO84227  
 Figure 79: DNA328369, BC007634, 90610.at  
 Figure 80A-B: DNA328370, NP\_001273.1, 200615.s.at  
 Figure 81: PRO84228  
 Figure 82: DNA323806, NP\_075385.1, 200644.at  
 Figure 83: PRO80555  
 Figure 84: DNA327532, GLUL, 200648.s.at  
 Figure 85: PRO71134  
 Figure 86: DNA227055, NP\_002625.1, 200658.s.at  
 Figure 87: PRO37518  
 Figure 88: DNA325702, NP\_001771.1, 200663.at  
 Figure 89: PRO283  
 Figure 90: DNA83172, NP\_003109.1, 200665.s.at  
 Figure 91: PRO2120  
 Figure 92: DNA328371, NP\_004347.1, 200675.at  
 Figure 93: PRO4866  
 Figure 94A-B: DNA328372, 105551.7, 200685.at  
 Figure 95: PRO84229  
 Figure 96: DNA324633, BC000478, 200691.s.at  
 Figure 97: PRO81277  
 Figure 98: DNA324633, NP\_004125.2, 200692.s.at  
 Figure 99: PRO81277  
 Figure 100: DNA88350, NP\_000168.1, 200696.s.at  
 Figure 101: PRO2758  
 Figure 102: DNA328373, AB034747, 200704.at  
 Figure 103: PRO84230  
 Figure 104: DNA328374, NP\_004853.1, 200706.s.at  
 Figure 105: PRO84231  
 Figure 106: DNA328375, NP\_002071.1, 200708.at  
 Figure 107: PRO80880

- Figure 108: DNA328376, NP\_001210.1, 200755.s.at  
 Figure 109: PRO1015  
 Figure 110A-B: DNA269826, NP\_003195.1, 200758.s.at  
 Figure 111: PRO58228  
 Figure 112: DNA325414, NP\_001900.1, 200766.at  
 Figure 113: PRO292  
 Figure 114A-C: DNA188738, NP\_002284.2, 200771.at  
 Figure 115: PRO25580  
 Figure 116: DNA328377, NP\_003759.1, 200787.s.at  
 Figure 117: PRO84232  
 Figure 118: DNA270954, NP\_001089.1, 200793.s.at  
 Figure 119: PRO59285  
 Figure 120: DNA272928, NP\_055579.1, 200794.x.at  
 Figure 121: PRO61012  
 Figure 122A-B: DNA327536, BC017197, 200797.s.at  
 Figure 123: PRO37003  
 Figure 124: DNA287211, NP\_002147.1, 200806.s.at  
 Figure 125: PRO69492  
 Figure 126: DNA326655, NP\_002803.1, 200820.at  
 Figure 127: PRO83005  
 Figure 128A-B: DNA328378, AB032261, 200832.s.at  
 Figure 129: PRO84233  
 Figure 130: DNA103558, NP\_005736.1, 200837.at  
 Figure 131: PRO4885  
 Figure 132: DNA196817, NP\_001899.1, 200838.at  
 Figure 133: PRO3344  
 Figure 134A-B: DNA327537, NP\_004437.1, 200842.s.at  
 Figure 135: PRO83581  
 Figure 136: DNA323982, NP\_004896.1, 200844.s.at  
 Figure 137: PRO80709  
 Figure 138: DNA323876, NP\_006612.2, 200850.s.at  
 Figure 139: PRO80619  
 Figure 140A-B: DNA228029, NP\_055577.1, 200862.at  
 Figure 141: PRO38492  
 Figure 142: DNA328379, BC015869, 200878.at  
 Figure 143: PRO84234  
 Figure 144: DNA325584, NP\_002005.1, 200895.s.at  
 Figure 145: PRO59262  
 Figure 146A-B: DNA274281, NP\_036347.1, 200899.s.at  
 Figure 147: PRO62204  
 Figure 148: DNA226028, NP\_002346.1, 200900.s.at  
 Figure 149: PRO36491  
 Figure 150: DNA326819, NP\_000678.1, 200903.s.at  
 Figure 151: PRO83152  
 Figure 152: DNA328380, HSHLAEHCM, 200904.at  
 Figure 153: DNA328381, NP\_005507.1, 200905.x.at  
 Figure 154: PRO84236  
 Figure 155: DNA272695, NP\_001722.1, 200920.s.at  
 Figure 156: PRO60817  
 Figure 157: DNA327255, NP\_002385.2, 200924.s.at  
 Figure 158: PRO57298  
 Figure 159: DNA327540, NP\_006818.1, 200929.at  
 Figure 160: PRO38005  
 Figure 161: DNA225878, NP\_004334.1, 200935.at  
 Figure 162: PRO36341  
 Figure 163: DNA328382, 160963.2, 200941.at  
 Figure 164: PRO84237  
 Figure 165: DNA328383, NP\_004956.3, 200944.s.at  
 Figure 166: PRO84238  
 Figure 167A-B: DNA287217, NP\_001750.1, 200953.s.at  
 Figure 168: PRO36766  
 Figure 169: DNA328384, NP\_036380.2, 200961.at  
 Figure 170: PRO84239  
 Figure 171: DNA328385, AK001310, 200972.at  
 Figure 172: PRO730  
 Figure 173: DNA326040, NP\_005715.1, 200973.s.at  
 Figure 174: PRO730  
 Figure 175: DNA324110, NP\_005908.1, 200978.at  
 Figure 176: PRO4918  
 Figure 177: DNA328386, NP\_000602.1, 200983.x.at  
 Figure 178: PRO2697  
 Figure 179: DNA275408, NP\_001596.1, 201000.at  
 Figure 180: PRO63068  
 Figure 181: DNA328387, NP\_001760.1, 201005.at  
 Figure 182: PRO4769  
 Figure 183: DNA103593, NP\_000174.1, 201007.at  
 Figure 184: PRO4917  
 Figure 185: DNA304713, NP\_006463.2, 201008.s.at  
 Figure 186: PRO71139  
 Figure 187: DNA328388, BC010273, 201013.s.at  
 Figure 188: PRO84240  
 Figure 189: DNA328389, NP\_006861.1, 201021.s.at  
 Figure 190: PRO84241  
 Figure 191: DNA328390, NP\_002291.1, 201030.x.at  
 Figure 192: PRO82116  
 Figure 193: DNA196628, NP\_005318.1, 201036.s.at  
 Figure 194: PRO25105  
 Figure 195: DNA287372, NP\_002618.1, 201037.at  
 Figure 196: PRO69632  
 Figure 197: DNA328391, NP\_004408.1, 201041.s.at  
 Figure 198: PRO84242  
 Figure 199: DNA196484, DNA196484, 201042.at  
 Figure 200: DNA227143, NP\_036400.1, 201050.at  
 Figure 201: PRO37606  
 Figure 202: DNA328392, 1500938.11, 201051.at  
 Figure 203: PRO84243  
 Figure 204: DNA328261, AF130103, 201060.x.at  
 Figure 205: DNA325001, NP\_002794.1, 201068.s.at  
 Figure 206: PRO81592  
 Figure 207: DNA328393, NP\_001651.1, 201096.s.at  
 Figure 208: PRO81010  
 Figure 209: DNA328394, AF131738, 201103.x.at  
 Figure 210A-B: DNA328395, NP\_056198.1, 201104.x.at  
 Figure 211: PRO84245  
 Figure 212: DNA328396, NP\_002076.1, 201106.at  
 Figure 213: PRO84246  
 Figure 214: DNA328397, NP\_002622.1, 201118.at



- Figure 215: PRO84247  
 Figure 216: DNA328398, NP\_002204.1, 201125.s.at  
 Figure 217: PRO34737  
 Figure 218: DNA325398, NP\_004083.2, 201135.at  
 Figure 219: PRO81930  
 Figure 220: DNA88520, NP\_002501.1, 201141.at  
 Figure 221: PRO2824  
 Figure 222: DNA324480, NP\_001544.1, 201163.s.at  
 Figure 223: PRO81141  
 Figure 224: DNA151802, NP\_003661.1, 201169.s.at  
 Figure 225: PRO12890  
 Figure 226: DNA226662, NP\_057043.1, 201175.at  
 Figure 227: PRO37125  
 Figure 228: DNA88066, NP\_002328.1, 201186.at  
 Figure 229: PRO2638  
 Figure 230: DNA273342, NP\_005887.1, 201193.at  
 Figure 231: PRO61345  
 Figure 232: DNA328399, NP\_003000.1, 201194.at  
 Figure 233: PRO84248  
 Figure 234A-B: DNA103453, HUME16GEN, 201195.s.at  
 Figure 235: PRO4780  
 Figure 236: DNA328400, NP\_003842.1, 201200.at  
 Figure 237: PRO1409  
 Figure 238: DNA327542, NP\_000091.1, 201201.at  
 Figure 239: PRO83582  
 Figure 240: DNA103488, NP\_002583.1, 201202.at  
 Figure 241: PRO4815  
 Figure 242: DNA328401, BC013678, 201212.at  
 Figure 243A-B: DNA328402, NP\_073713.1, 201220.x.at  
 Figure 244: PRO84249  
 Figure 245: DNA325380, NP\_004995.1, 201226.at  
 Figure 246: PRO81914  
 Figure 247: DNA226615, NP\_001668.1, 201242.s.at  
 Figure 248: PRO37078  
 Figure 249: DNA328403, NP\_037462.1, 201243.s.at  
 Figure 250: PRO84250  
 Figure 251: DNA270950, NP\_003182.1, 201263.at  
 Figure 252: PRO59281  
 Figure 253A-B: DNA328404, NP\_003321.1, 201266.at  
 Figure 254: PRO84251  
 Figure 255: DNA97290, NP\_002503.1, 201268.at  
 Figure 256: PRO3637  
 Figure 257: DNA325028, NP\_001619.1, 201272.at  
 Figure 258: PRO81617  
 Figure 259: DNA328405, NP\_112556.1, 201277.s.at  
 Figure 260: PRO84252  
 Figure 261: DNA328406, NP\_001334.1, 201279.s.at  
 Figure 262: PRO84253  
 Figure 263: DNA328407, WSB1, 201296.s.at  
 Figure 264: PRO84254  
 Figure 265: DNA328408, NP\_060713.1, 201308.s.at  
 Figure 266: PRO84255  
 Figure 267: DNA325595, NP\_001966.1, 201313.at  
 Figure 268: PRO38010  
 Figure 269: DNA255078, NP\_006426.1, 201315.x.at  
 Figure 270: PRO50165  
 Figure 271: DNA150781, NP\_001414.1, 201324.at  
 Figure 272: PRO12467  
 Figure 273: DNA328409, NP\_002075.2, 201348.at  
 Figure 274: PRO81281  
 Figure 275: DNA324475, NP\_004172.2, 201387.s.at  
 Figure 276: PRO81137  
 Figure 277: DNA226353, NP\_005769.1, 201395.at  
 Figure 278: PRO36816  
 Figure 279: DNA328410, NP\_004519.1, 201403.s.at  
 Figure 280: PRO60174  
 Figure 281A-B: DNA328411, 1400253.2, 201408.at  
 Figure 282: PRO84256  
 Figure 283: DNA328412, NP\_060428.1, 201411.s.at  
 Figure 284: PRO84257  
 Figure 285: DNA273517, NP\_000169.1, 201415.at  
 Figure 286: PRO61498  
 Figure 287: DNA327550, NP\_001959.1, 201435.s.at  
 Figure 288: PRO81164  
 Figure 289: DNA273396, DNA273396, 201449.at  
 Figure 290: DNA325049, NP\_005605.1, 201453.x.at  
 Figure 291: PRO37938  
 Figure 292: DNA274343, NP\_000894.1, 201467.s.at  
 Figure 293: PRO62259  
 Figure 294: DNA328413, NP\_004823.1, 201470.at  
 Figure 295: PRO84258  
 Figure 296: DNA328414, NP\_003891.1, 201471.s.at  
 Figure 297: PRO81346  
 Figure 298: DNA103320, NP\_002220.1, 201473.at  
 Figure 299: PRO4650  
 Figure 300: DNA88608, NP\_002893.1, 201485.s.at  
 Figure 301: PRO2864  
 Figure 302: DNA304459, BC005020, 201489.at  
 Figure 303: PRO37073  
 Figure 304: DNA304459, NP\_005720.1, 201490.s.at  
 Figure 305: PRO37073  
 Figure 306: DNA253807, NP\_065390.1, 201502.s.at  
 Figure 307: PRO49210  
 Figure 308: DNA328415, BC006997, 201503.at  
 Figure 309: PRO60207  
 Figure 310: DNA328416, NP\_002613.2, 201507.at  
 Figure 311: PRO84259  
 Figure 312: DNA271931, NP\_005745.1, 201514.s.at  
 Figure 313: PRO60207  
 Figure 314A-B: DNA150463, NP\_055635.1, 201519.at  
 Figure 315: PRO12269  
 Figure 316: DNA328417, ATP6V1F, 201527.at  
 Figure 317: PRO84260  
 Figure 318: DNA328418, NP\_003398.1, 201531.at  
 Figure 319: PRO84261  
 Figure 320: DNA328419, NP\_002779.1, 201532.at  
 Figure 321: PRO84262  
 Figure 322: DNA328420, BC002682, 201537.s.at  
 Figure 323: PRO58245  
 Figure 324: DNA88464, NP\_005552.2, 201551.s.at

- Figure 325: PRO2804  
 Figure 326A-B: DNA290226, NP\_039234.1, 201559.s.at  
 Figure 327: PRO70317  
 Figure 328: DNA227071, NP\_000260.1, 201577.at  
 Figure 329: PRO37534  
 Figure 330A-B: DNA227307, NP\_009115.1, 201591.s.at  
 Figure 331: PRO37770  
 Figure 332: DNA255406, NP\_005533.1, 201625.s.at  
 Figure 333: PRO50473  
 Figure 334A-B: DNA328421, 475621.10, 201646.at  
 Figure 335: PRO51048  
 Figure 336A-B: DNA220748, NP\_000201.1, 201656.at  
 Figure 337: PRO34726  
 Figure 338: DNA269791, NP\_001168.1, 201659.s.at  
 Figure 339: PRO58197  
 Figure 340A-B: DNA328422, NP\_004448.1, 201661.s.at  
 Figure 341: PRO84263  
 Figure 342: DNA328423, NP\_003245.1, 201666.at  
 Figure 343: PRO2121  
 Figure 344: DNA273090, NP\_002347.4, 201670.s.at  
 Figure 345: PRO61148  
 Figure 346: DNA328424, NP\_005142.1, 201672.s.at  
 Figure 347: PRO59291  
 Figure 348: DNA271223, NP\_005070.1, 201689.s.at  
 Figure 349: PRO59538  
 Figure 350A-B: DNA323965, NP\_002848.1, 201706.s.at  
 Figure 351: PRO80695  
 Figure 352: DNA270883, NP\_001061.1, 201714.at  
 Figure 353: PRO59218  
 Figure 354A-B: DNA328425, NP\_065207.2, 201722.s.at  
 Figure 355: PRO84264  
 Figure 356: DNA328426, NP\_000582.1, 201743.at  
 Figure 357: PRO384  
 Figure 358: DNA150429, NP\_002813.1, 201745.at  
 Figure 359: PRO12769  
 Figure 360: DNA272465, NP\_004543.1, 201757.at  
 Figure 361: PRO60713  
 Figure 362: DNA328427, NP\_061109.1, 201760.s.at  
 Figure 363: PRO84265  
 Figure 364: DNA287167, NP\_006627.1, 201761.at  
 Figure 365: PRO59136  
 Figure 366: DNA323937, NP\_005689.2, 201771.at  
 Figure 367: PRO80670  
 Figure 368: DNA88619, NP\_002924.1, 201785.at  
 Figure 369: PRO2871  
 Figure 370A-B: DNA328428, NP\_038479.1, 201798.s.at  
 Figure 371: PRO84266  
 Figure 372: DNA227563, NP\_004946.1, 201801.s.at  
 Figure 373: PRO38026  
 Figure 374: DNA225896, NP\_000109.1, 201808.s.at  
 Figure 375: PRO36359  
 Figure 376: DNA151017, NP\_004835.1, 201810.s.at  
 Figure 377: PRO12841  
 Figure 378: DNA328429, NP\_079106.2, 201818.at  
 Figure 379: PRO81201  
 Figure 380: DNA328430, NP\_005496.2, 201819.at  
 Figure 381: PRO84267  
 Figure 382: DNA324015, NP\_006326.1, 201821.s.at  
 Figure 383: PRO80735  
 Figure 384: DNA150650, NP\_057086.1, 201825.s.at  
 Figure 385: PRO12393  
 Figure 386: DNA304710, NP\_001531.1, 201841.s.at  
 Figure 387: PRO71136  
 Figure 388: DNA88450, NP\_000226.1, 201847.at  
 Figure 389: PRO2795  
 Figure 390: DNA150725, NP\_001738.1, 201850.at  
 Figure 391: PRO12792  
 Figure 392: DNA272066, NP\_002931.1, 201872.s.at  
 Figure 393: PRO60337  
 Figure 394: DNA328431, NP\_001817.1, 201897.s.at  
 Figure 395: PRO45093  
 Figure 396: DNA103214, NP\_006057.1, 201900.s.at  
 Figure 397: PRO4544  
 Figure 398: DNA227112, NP\_006397.1, 201923.at  
 Figure 399: PRO37575  
 Figure 400: DNA83046, NP\_000565.1, 201926.s.at  
 Figure 401: PRO2569  
 Figure 402: DNA273014, NP\_000117.1, 201931.at  
 Figure 403: PRO61085  
 Figure 404: DNA254147, NP\_000512.1, 201944.at  
 Figure 405: PRO49262  
 Figure 406: DNA274167, NP\_006422.1, 201946.s.at  
 Figure 407: PRO62097  
 Figure 408A-B: DNA327562, HSMEMD, 201951.at  
 Figure 409A-B: DNA327563, NP\_066945.1, 201963.at  
 Figure 410: PRO83592  
 Figure 411: DNA227290, NP\_055861.1, 201965.s.at  
 Figure 412: PRO37753  
 Figure 413A-B: DNA328432, NP\_005768.1, 201967.at  
 Figure 414: PRO61793  
 Figure 415A-B: DNA328433, ATP6V1A1, 201971.s.at  
 Figure 416: PRO84268  
 Figure 417: DNA327073, NP\_036418.1, 201994.at  
 Figure 418: PRO83365  
 Figure 419: DNA226878, NP\_000118.1, 201995.at  
 Figure 420: PRO37341  
 Figure 421A-D: DNA328434, NP\_055816.2, 201996.s.at  
 Figure 422: PRO84269  
 Figure 423: DNA328435, NP\_002481.1, 202001.s.at  
 Figure 424: PRO60236  
 Figure 425: DNA275246, NP\_006102.1, 202003.s.at  
 Figure 426: PRO62933  
 Figure 427: DNA327841, NP\_068813.1, 202005.at  
 Figure 428: PRO12377

- Figure 429: DNA328436, 1171619.4, 202007.at  
 Figure 430: PRO84270  
 Figure 431: DNA327564, NP\_000111.1, 202017.at  
 Figure 432: PRO83593  
 Figure 433: DNA328437, AF083441, 202021.x.at  
 Figure 434: PRO84271  
 Figure 435A-B: DNA270997, NP\_005047.1, 202040.s.at  
 Figure 436: PRO59326  
 Figure 437A-B: DNA327565, NP\_056392.1, 202052.s.at  
 Figure 438: PRO83594  
 Figure 439A-B: DNA327566, NP\_000373.1, 202053.s.at  
 Figure 440: PRO83595  
 Figure 441: DNA226116, NP\_002990.1, 202071.at  
 Figure 442: PRO36579  
 Figure 443A-B: DNA328438, 100983.30, 202073.at  
 Figure 444: PRO84272  
 Figure 445: DNA328439, NP\_068815.1, 202074.s.at  
 Figure 446: PRO84273  
 Figure 447: DNA290272, NP\_004898.1, 202081.at  
 Figure 448: PRO70409  
 Figure 449: DNA327569, NP\_001903.1, 202087.s.at  
 Figure 450: PRO2683  
 Figure 451: DNA328440, NP\_004517.1, 202107.s.at  
 Figure 452: PRO84274  
 Figure 453: DNA272777, NP\_000276.1, 202108.at  
 Figure 454: PRO60884  
 Figure 455A-B: DNA328441, AL136139, 202149.at  
 Figure 456: PRO0  
 Figure 457: DNA328442, NP\_006078.2, 202154.x.at  
 Figure 458: PRO84275  
 Figure 459A-C: DNA328443, NP\_004371.1, 202160.at  
 Figure 460: PRO84276  
 Figure 461A-C: DNA271201, NP\_005881.1, 202191.s.at  
 Figure 462: PRO59518  
 Figure 463: DNA328258, SLC16A1, 202236.s.at  
 Figure 464: PRO84151  
 Figure 465: DNA328444, MGC14458, 202246.s.at  
 Figure 466: PRO84277  
 Figure 467: DNA294794, NP\_002861.1, 202252.at  
 Figure 468: PRO70754  
 Figure 469A-B: DNA227176, NP\_056371.1, 202255.s.at  
 Figure 470: PRO37639  
 Figure 471: DNA325823, NP\_055702.1, 202258.s.at  
 Figure 472: PRO82289  
 Figure 473: DNA256533, NP\_006105.1, 202264.s.at  
 Figure 474: PRO51565  
 Figure 475: DNA328445, NP\_057698.1, 202266.at  
 Figure 476: PRO84278  
 Figure 477: DNA328446, NP\_003896.1, 202268.s.at  
 Figure 478: PRO59821  
 Figure 479: DNA328447, NP\_000393.2, 202275.at  
 Figure 480: PRO84279  
 Figure 481: DNA304716, NP\_510867.1, 202284.s.at  
 Figure 482: PRO71142  
 Figure 483: DNA270142, NP\_005947.2, 202309.at  
 Figure 484: PRO58531  
 Figure 485: DNA328448, NP\_000777.1, 202314.at  
 Figure 486: PRO62362  
 Figure 487: DNA325115, NP\_001435.1, 202345.s.at  
 Figure 488: PRO81689  
 Figure 489: DNA106239, DNA106239, 202351.at  
 Figure 490: DNA270502, NP\_002807.1, 202352.s.at  
 Figure 491: PRO58880  
 Figure 492: DNA327074, FLJ21174, 202371.at  
 Figure 493: PRO83366  
 Figure 494: DNA149091, DNA149091, 202377.at  
 Figure 495A-B: DNA151045, NP\_005376.2, 202379.s.at  
 Figure 496: PRO12587  
 Figure 497A-B: DNA200236, NP\_003807.1, 202381.at  
 Figure 498: PRO34137  
 Figure 499: DNA328449, NP\_005462.1, 202382.s.at  
 Figure 500: PRO60304  
 Figure 501: DNA290234, NP\_002914.1, 202388.at  
 Figure 502: PRO70333  
 Figure 503: DNA269766, NP\_005646.1, 202393.s.at  
 Figure 504: PRO58175  
 Figure 505: DNA227612, NP\_056230.1, 202427.s.at  
 Figure 506: PRO38075  
 Figure 507: DNA324171, NP\_065438.1, 202428.x.at  
 Figure 508: PRO60753  
 Figure 509A-B: DNA327576, NP\_000095.1, 202434.s.at  
 Figure 510: PRO83600  
 Figure 511A-D: DNA328450, NP\_077719.1, 202443.x.at  
 Figure 512: PRO84280  
 Figure 513: DNA225809, NP\_000387.1, 202450.s.at  
 Figure 514: PRO36272  
 Figure 515: DNA227921, NP\_003789.1, 202468.s.at  
 Figure 516: PRO38384  
 Figure 517: DNA150942, HSY18007, 202475.at  
 Figure 518: PRO12549  
 Figure 519: DNA225566, NP\_004744.1, 202481.at  
 Figure 520: PRO36029  
 Figure 521A-B: DNA103449, NP\_008862.1, 202497.x.at  
 Figure 522: PRO4776  
 Figure 523: DNA328451, NP\_000007.1, 202502.at  
 Figure 524: PRO62139  
 Figure 525A-B: DNA274893, NP\_006282.1, 202510.s.at  
 Figure 526: PRO62634  
 Figure 527: DNA328452, NP\_000394.1, 202528.at  
 Figure 528: PRO63289  
 Figure 529: DNA219229, NP\_002189.1, 202531.at  
 Figure 530: PRO34544

Figure 531A-B: DNA274852, NP\_004115.1, 202543.s\_at  
 Figure 532: PRO62605  
 Figure 533: DNA328453, NP\_003752.2, 202546.at  
 Figure 534: PRO84281  
 Figure 535A-B: DNA328454, NP\_057525.1, 202551.s\_at  
 Figure 536: PRO4330  
 Figure 537: DNA150817, NP\_000840.1, 202554.s\_at  
 Figure 538: PRO12808  
 Figure 539: DNA227994, NP\_009107.1, 202562.s\_at  
 Figure 540: PRO38457  
 Figure 541: DNA328455, AY007134, 202573.at  
 Figure 542: PRO84282  
 Figure 543: DNA323923, NP\_001869.1, 202575.at  
 Figure 544: PRO80657  
 Figure 545: DNA328456, NP\_000467.1, 202587.s\_at  
 Figure 546: PRO84283  
 Figure 547: DNA328457, NP\_036422.1, 202606.s\_at  
 Figure 548: PRO70421  
 Figure 549: DNA103245, NP\_002341.1, 202626.s\_at  
 Figure 550: PRO4575  
 Figure 551: DNA83141, NP\_000593.1, 202627.s\_at  
 Figure 552: PRO2604  
 Figure 553: DNA254129, NP\_006001.1, 202655.at  
 Figure 554: PRO49244  
 Figure 555: DNA270379, NP\_002792.1, 202659.at  
 Figure 556: PRO58763  
 Figure 557: DNA326896, NP\_003672.1, 202671.s\_at  
 Figure 558: PRO69486  
 Figure 559: DNA289526, NP\_004015.2, 202672.s\_at  
 Figure 560: PRO70282  
 Figure 561: DNA273542, NP\_002991.1, 202675.at  
 Figure 562: PRO61522  
 Figure 563: DNA328458, NP\_037458.2, 202679.at  
 Figure 564: PRO84284  
 Figure 565: DNA84130, NP\_003801.1, 202687.s\_at  
 Figure 566: PRO1096  
 Figure 567: DNA271085, NP\_004751.1, 202693.s\_at  
 Figure 568: PRO59409  
 Figure 569A-B: DNA150467, NP\_055513.1, 202699.s\_at  
 Figure 570: PRO12272  
 Figure 571A-B: DNA328459, NP\_004332.2, 202715.at  
 Figure 572: PRO84285  
 Figure 573: DNA273290, NP\_002047.1, 202722.s\_at  
 Figure 574: PRO61300  
 Figure 575: DNA328460, NP\_004190.1, 202733.at  
 Figure 576: PRO84286  
 Figure 577: DNA150713, NP\_006570.1, 202735.at  
 Figure 578: PRO12082  
 Figure 579A-B: DNA328461, 350230.2, 202741.at  
 Figure 580: PRO84287  
 Figure 581: DNA271973, NP\_002722.1, 202742.s\_at  
 Figure 582: PRO60248  
 Figure 583A-B: DNA150943, NP\_036376.1, 202752.x\_at  
 Figure 584: PRO12550  
 Figure 585A-C: DNA328462, HSA303079, 202759.s\_at  
 Figure 586: PRO84288  
 Figure 587A-C: DNA328463, NP\_009134.1, 202760.s\_at  
 Figure 588: PRO84289  
 Figure 589: DNA226080, NP\_001601.1, 202767.at  
 Figure 590: PRO36543  
 Figure 591A-B: DNA150977, NP\_006723.1, 202768.at  
 Figure 592: PRO12828  
 Figure 593A-B: DNA328464, 977954.20, 202769.at  
 Figure 594: PRO84290  
 Figure 595: DNA226578, NP\_004345.1, 202770.s\_at  
 Figure 596: PRO37041  
 Figure 597A-B: DNA103521, NP\_004163.1, 202800.at  
 Figure 598: PRO4848  
 Figure 599A-B: DNA327583, ABCC1, 202805.s\_at  
 Figure 600: PRO83604  
 Figure 601: DNA328465, NP\_005639.1, 202823.at  
 Figure 602: PRO84291  
 Figure 603: DNA225865, NP\_004986.1, 202827.s\_at  
 Figure 604: PRO36328  
 Figure 605: DNA225926, NP\_000138.1, 202838.at  
 Figure 606: PRO36389  
 Figure 607: DNA328466, NP\_004554.1, 202847.at  
 Figure 608: PRO84292  
 Figure 609: DNA103394, NP\_004198.1, 202855.s\_at  
 Figure 610: PRO4722  
 Figure 611: DNA275144, NP\_000128.1, 202862.at  
 Figure 612: PRO62852  
 Figure 613: DNA328467, SP100, 202864.s\_at  
 Figure 614: PRO84293  
 Figure 615: DNA287289, NP\_058132.1, 202869.at  
 Figure 616: PRO69559  
 Figure 617: DNA328468, BC010960, 202872.at  
 Figure 618: PRO84294  
 Figure 619: DNA328469, NP\_001686.1, 202874.s\_at  
 Figure 620: PRO84295  
 Figure 621A-B: DNA255318, NP\_036204.1, 202877.s\_at  
 Figure 622: PRO50388  
 Figure 623A-B: DNA328470, NP\_055620.1, 202909.at  
 Figure 624: PRO84296  
 Figure 625: DNA327584, NP\_002955.2, 202917.s\_at  
 Figure 626: PRO80649  
 Figure 627: DNA272425, NP\_001489.1, 202923.s\_at  
 Figure 628: PRO60677  
 Figure 629: DNA328471, ZMPSTE24, 202939.at  
 Figure 630: PRO84297  
 Figure 631: DNA269481, NP\_001976.1, 202942.at  
 Figure 632: PRO57901  
 Figure 633: DNA328472, NP\_000482.2, 202953.at  
 Figure 634: PRO84298  
 Figure 635A-B: DNA328473, NP\_006473.1,

- 202968.s.at  
Figure 636: PRO84299  
Figure 637A-C: DNA328474, 1501914.1, 202969.at  
Figure 638: PRO84300  
Figure 639: DNA325915, ZAP128, 202982.s.at  
Figure 640: PRO82369  
Figure 641: DNA271272, NP\_000366.1, 203031.s.at  
Figure 642: PRO59583  
Figure 643: DNA324049, FH, 203032.s.at  
Figure 644: PRO62607  
Figure 645A-B: DNA271865, NP\_055566.1, 203037.s.at  
Figure 646: PRO60145  
Figure 647: DNA328475, LAMP2, 203042.at  
Figure 648: PRO84301  
Figure 649A-B: DNA328476, AF074331, 203058.s.at  
Figure 650: PRO84302  
Figure 651: DNA256830, NP\_004815.1, 203100.s.at  
Figure 652: PRO51761  
Figure 653: DNA272867, NP\_003960.1, 203109.at  
Figure 654: PRO60960  
Figure 655A-B: DNA227582, NP\_000608.1, 203124.s.at  
Figure 656: PRO38045  
Figure 657: DNA328477, NP\_003767.1, 203152.at  
Figure 658: PRO84303  
Figure 659A-B: DNA328478, NP\_055720.2, 203158.s.at  
Figure 660: PRO84304  
Figure 661: DNA226136, NP\_003246.1, 203167.at  
Figure 662: PRO36599  
Figure 663: DNA328479, NP\_001473.1, 203178.at  
Figure 664: PRO84305  
Figure 665A-C: DNA328480, NP\_001990.1, 203184.at  
Figure 666: PRO84306  
Figure 667A-B: DNA271010, NP\_055552.1, 203185.at  
Figure 668: PRO59339  
Figure 669: DNA270448, NP\_002487.1, 203189.s.at  
Figure 670: PRO58827  
Figure 671A-B: DNA328481, MTMR2, 203211.s.at  
Figure 672: PRO84307  
Figure 673A-C: DNA328482, NP\_000426.1, 203238.s.at  
Figure 674: PRO84308  
Figure 675: DNA328483, NP\_061163.1, 203255.at  
Figure 676: PRO84309  
Figure 677: DNA227127, NP\_003571.1, 203269.at  
Figure 678: PRO37590  
Figure 679: DNA328484, UNC119, 203271.s.at  
Figure 680: PRO84310  
Figure 681: DNA302020, NP\_005564.1, 203276.at  
Figure 682: PRO70993  
Figure 683A-B: DNA328485, BHC80, 203278.s.at  
Figure 684: PRO84311  
Figure 685: DNA328486, NP\_000149.1, 203282.at  
Figure 686: PRO60119  
Figure 687: DNA328487, AF251295, 203299.s.at  
Figure 688: PRO84312  
Figure 689: DNA328488, NP\_003907.2, 203300.x.at  
Figure 690: PRO84313  
Figure 691: DNA328489, NP\_006511.1, 203303.at  
Figure 692: PRO84314  
Figure 693A-B: DNA328490, NP\_000120.1, 203305.at  
Figure 694: PRO84315  
Figure 695: DNA327593, NP\_006205.1, 203335.at  
Figure 696: PRO59733  
Figure 697: DNA328491, ICAP-1A, 203336.s.at  
Figure 698: PRO61323  
Figure 699A-B: DNA328492, NP\_056125.1, 203354.s.at  
Figure 700: PRO84316  
Figure 701: DNA328493, NP\_008957.1, 203367.at  
Figure 702: PRO84317  
Figure 703: DNA328494, RPS6KA1, 203379.at  
Figure 704: PRO84318  
Figure 705: DNA274960, NP\_008856.1, 203380.x.at  
Figure 706: PRO62694  
Figure 707: DNA88084, NP\_000032.1, 203381.s.at  
Figure 708: PRO2644  
Figure 709A-B: DNA254616, NP\_004473.1, 203397.s.at  
Figure 710: PRO49718  
Figure 711: DNA326892, NP\_003711.1, 203405.at  
Figure 712: PRO83213  
Figure 713: DNA323927, NP\_005563.1, 203411.s.at  
Figure 714: PRO80660  
Figure 715: DNA151037, NP\_036461.1, 203414.at  
Figure 716: PRO12586  
Figure 717: DNA273410, NP\_004036.1, 203454.s.at  
Figure 718: PRO61409  
Figure 719: DNA328495, NP\_055578.1, 203465.at  
Figure 720: PRO58967  
Figure 721: DNA328496, NP\_002428.1, 203466.at  
Figure 722: PRO80786  
Figure 723A-B: DNA255622, NP\_009187.1, 203472.s.at  
Figure 724: PRO50686  
Figure 725A-C: DNA328497, NP\_005493.1, 203504.s.at  
Figure 726: PRO84319  
Figure 727A-C: DNA328498, AF285167, 203505.at  
Figure 728: PRO84320  
Figure 729A-B: DNA188400, NP\_001057.1, 203508.at  
Figure 730: PRO21928  
Figure 731A-B: DNA328499, NP\_003096.1, 203509.at  
Figure 732: PRO84321  
Figure 733: DNA272911, NP\_006545.1, 203517.at  
Figure 734: PRO60997  
Figure 735A-D: DNA328500, NP\_000072.1, 203518.at  
Figure 736: PRO84322  
Figure 737A-B: DNA103296, NP\_006369.1, 203528.at

- Figure 738: PRO4626  
 Figure 739: DNA323910, NP\_002956.1, 203535.at  
 Figure 740: PRO80648  
 Figure 741A-B: DNA272399, NP\_001197.1, 203543.s.at  
 Figure 742: PRO60653  
 Figure 743: DNA328501, NP\_076984.1, 203545.at  
 Figure 744: PRO84323  
 Figure 745: DNA88453, NP\_000228.1, 203548.s.at  
 Figure 746: PRO2797  
 Figure 747: DNA328502, NP\_006566.2, 203553.s.at  
 Figure 748: PRO84324  
 Figure 749: DNA328503, NP\_000272.1, 203557.s.at  
 Figure 750: PRO10850  
 Figure 751: DNA327594, NP\_003869.1, 203560.at  
 Figure 752: PRO83611  
 Figure 753: DNA225916, NP\_067674.1, 203561.at  
 Figure 754: PRO36379  
 Figure 755: DNA273676, NP\_055488.1, 203584.at  
 Figure 756: PRO61644  
 Figure 757: DNA83085, NP\_000751.1, 203591.s.at  
 Figure 758: PRO2583  
 Figure 759: DNA271003, NP\_003720.1, 203594.at  
 Figure 760: PRO59332  
 Figure 761A-B: DNA328504, 1400155.1, 203608.at  
 Figure 762: PRO84325  
 Figure 763: DNA328505, NP\_002484.1, 203613.s.at  
 Figure 764: PRO62117  
 Figure 765: DNA328506, NP\_001046.1, 203615.x.at  
 Figure 766: PRO84326  
 Figure 767: DNA225774, NP\_005079.1, 203624.at  
 Figure 768: PRO36237  
 Figure 769: DNA254642, NP\_004100.1, 203646.at  
 Figure 770: PRO49743  
 Figure 771: DNA328507, NP\_006395.1, 203650.at  
 Figure 772: PRO4761  
 Figure 773A-B: DNA272998, NP\_055548.1, 203651.at  
 Figure 774: PRO61070  
 Figure 775: DNA328508, NP\_003368.1, 203683.s.at  
 Figure 776: PRO35975  
 Figure 777: DNA255298, NP\_004394.1, 203695.s.at  
 Figure 778: PRO50371  
 Figure 779: DNA227020, NP\_001416.1, 203729.at  
 Figure 780: PRO37483  
 Figure 781: DNA328509, NP\_006739.1, 203760.s.at  
 Figure 782: PRO57996  
 Figure 783: DNA328510, NP\_055066.1, 203775.at  
 Figure 784: PRO84327  
 Figure 785A-B: DNA194602, NP\_006370.1, 203789.s.at  
 Figure 786: PRO23944  
 Figure 787: DNA328511, NP\_031397.1, 203825.at  
 Figure 788: PRO57838  
 Figure 789A-B: DNA328512, NP\_005772.2, 203839.s.at  
 Figure 790: PRO84328  
 Figure 791A-B: DNA272451, HSU86453, 203879.at  
 Figure 792: PRO60700  
 Figure 793: DNA82429, NP\_003011.1, 203889.at  
 Figure 794: PRO2558  
 Figure 795: DNA328513, NP\_057367.1, 203893.at  
 Figure 796: PRO37815  
 Figure 797: DNA150974, NP\_005684.1, 203920.at  
 Figure 798: PRO12224  
 Figure 799: DNA271676, NP\_002052.1, 203925.at  
 Figure 800: PRO59961  
 Figure 801: DNA88239, NP\_004985.1, 203936.s.at  
 Figure 802: PRO2711  
 Figure 803: DNA227232, NP\_001850.1, 203971.at  
 Figure 804: PRO37695  
 Figure 805: DNA328514, NP\_005186.1, 203973.s.at  
 Figure 806: PRO84329  
 Figure 807: DNA328515, NP\_000775.1, 203979.at  
 Figure 808: PRO84330  
 Figure 809: DNA327608, NP\_001433.1, 203980.at  
 Figure 810: PRO83617  
 Figure 811: DNA328516, NP\_005833.1, 204011.at  
 Figure 812: PRO12323  
 Figure 813: DNA328517, NP\_003558.1, 204032.at  
 Figure 814: PRO84331  
 Figure 815: DNA226342, NP\_000305.1, 204054.at  
 Figure 816: PRO36805  
 Figure 817: DNA327609, 1448428.2, 204058.at  
 Figure 818: PRO83618  
 Figure 819: DNA328518, ME1, 204059.s.at  
 Figure 820: PRO84332  
 Figure 821: DNA226737, NP\_004576.1, 204070.at  
 Figure 822: PRO37200  
 Figure 823A-C: DNA328519, NP\_075463.1, 204072.s.at  
 Figure 824: PRO84333  
 Figure 825: DNA328520, NP\_079353.1, 204080.at  
 Figure 826: PRO84334  
 Figure 827A-B: DNA150739, NP\_006484.1, 204084.s.at  
 Figure 828: PRO12442  
 Figure 829: DNA227130, NP\_002551.1, 204088.at  
 Figure 830: PRO37593  
 Figure 831: DNA328521, NP\_003069.1, 204099.at  
 Figure 832: PRO62553  
 Figure 833: DNA328522, NP\_001769.2, 204118.at  
 Figure 834: PRO2696  
 Figure 835: DNA328523, NP\_006712.1, 204119.s.at  
 Figure 836: PRO84335  
 Figure 837: DNA328524, NP\_057097.1, 204125.at  
 Figure 838: PRO84336  
 Figure 839: DNA328525, BC021224, 204131.s.at  
 Figure 840: PRO84337  
 Figure 841: DNA103532, NP\_003263.1, 204137.at  
 Figure 842: PRO4859  
 Figure 843: DNA324816, NP\_001060.1, 204141.at  
 Figure 844: PRO81429

- Figure 845: DNA270524, NP\_059982.1, 204142.at  
 Figure 846: PRO58901  
 Figure 847: DNA328526, NP\_000841.1, 204149.s.at  
 Figure 848: PRO37856  
 Figure 849A-B: DNA150497, DNA150497, 204155.s.at  
 Figure 850: PRO12296  
 Figure 851A-B: DNA328527, NP\_055751.1, 204160.s.at  
 Figure 852: PRO4351  
 Figure 853: DNA328528, MLC1SA, 204173.at  
 Figure 854: PRO60636  
 Figure 855: DNA328529, NP\_001620.2, 204174.at  
 Figure 856: PRO49814  
 Figure 857: DNA226380, NP\_001765.1, 204192.at  
 Figure 858: PRO4695  
 Figure 859: DNA273070, NP\_005189.2, 204193.at  
 Figure 860: PRO70107  
 Figure 861: DNA227514, NP\_000152.1, 204224.s.at  
 Figure 862: PRO37977  
 Figure 863: DNA270434, NP\_006434.1, 204238.s.at  
 Figure 864: PRO58814  
 Figure 865: DNA307936, NP\_004926.1, 204247.s.at  
 Figure 866: PRO71356  
 Figure 867A-B: DNA188734, NP\_001261.1, 204258.at  
 Figure 868: PRO22296  
 Figure 869: DNA226577, NP\_071390.1, 204265.s.at  
 Figure 870: PRO37040  
 Figure 871: DNA273802, NP\_066950.1, 204285.s.at  
 Figure 872: PRO61763  
 Figure 873: DNA328530, NP\_009198.2, 204328.at  
 Figure 874: PRO24118  
 Figure 875: DNA328531, NP\_037542.1, 204348.s.at  
 Figure 876: PRO84338  
 Figure 877: DNA328532, LIMK1, 204357.s.at  
 Figure 878: PRO84339  
 Figure 879: DNA225750, NP\_000254.1, 204360.s.at  
 Figure 880: PRO36213  
 Figure 881: DNA328533, NP\_003647.1, 204392.at  
 Figure 882: PRO84340  
 Figure 883: DNA272469, NP\_005299.1, 204396.s.at  
 Figure 884: PRO60717  
 Figure 885: DNA226462, NP\_002241.1, 204401.at  
 Figure 886: PRO36925  
 Figure 887: DNA225756, NP\_001636.1, 204416.x.at  
 Figure 888: PRO36219  
 Figure 889: DNA226286, NP\_001657.1, 204425.at  
 Figure 890: PRO36749  
 Figure 891A-B: DNA88476, NP\_002429.1, 204438.at  
 Figure 892: PRO2811  
 Figure 893: DNA150972, NP\_005252.1, 204472.at  
 Figure 894: PRO12162  
 Figure 895: DNA194652, NP\_001187.1, 204493.at  
 Figure 896: PRO23974  
 Figure 897: DNA328534, NP\_056307.1, 204494.s.at  
 Figure 898: PRO84341  
 Figure 899: DNA328254, BC002678, 204517.at  
 Figure 900: PRO11581  
 Figure 901: DNA328254, NP\_000934.1, 204518.s.at  
 Figure 902: PRO11581  
 Figure 903A-B: DNA328535, NP\_009147.1, 204544.at  
 Figure 904: PRO60044  
 Figure 905: DNA225993, NP\_000646.1, 204563.at  
 Figure 906: PRO36456  
 Figure 907: DNA287284, NP\_060943.1, 204565.at  
 Figure 908: PRO59915  
 Figure 909: DNA151910, NP\_004906.2, 204567.s.at  
 Figure 910: PRO12754  
 Figure 911: DNA270564, NP\_004499.1, 204615.x.at  
 Figure 912: PRO58939  
 Figure 913: DNA328536, 1099945.20, 204619.s.at  
 Figure 914: PRO84342  
 Figure 915A-D: DNA328537, NP\_004376.2, 204620.s.at  
 Figure 916: PRO84343  
 Figure 917: DNA151048, NP\_006177.1, 204621.s.at  
 Figure 918: PRO12850  
 Figure 919A-B: DNA328538, 351122.2, 204627.s.at  
 Figure 920: PRO84344  
 Figure 921A-B: DNA88429, NP\_000203.1, 204628.s.at  
 Figure 922: PRO2344  
 Figure 923: DNA226079, NP\_001602.1, 204638.at  
 Figure 924: PRO36542  
 Figure 925: DNA272078, NP\_003019.1, 204657.s.at  
 Figure 926: PRO60348  
 Figure 927: DNA227425, NP\_001038.1, 204675.at  
 Figure 928: PRO37888  
 Figure 929A-B: DNA328539, NP\_000121.1, 204713.s.at  
 Figure 930: PRO84345  
 Figure 931: DNA328540, NP\_006144.1, 204725.s.at  
 Figure 932: PRO12168  
 Figure 933A-B: DNA325192, NP\_038203.1, 204744.s.at  
 Figure 934: PRO81753  
 Figure 935: DNA328541, NP\_004503.1, 204773.at  
 Figure 936: PRO4843  
 Figure 937: DNA328542, NP\_055025.1, 204774.at  
 Figure 938: PRO2577  
 Figure 939: DNA327050, NP\_009199.1, 204787.at  
 Figure 940: PRO34043  
 Figure 941: DNA328543, NP\_005883.1, 204789.at  
 Figure 942: PRO84346  
 Figure 943: DNA272121, NP\_005895.1, 204790.at  
 Figure 944: PRO60391  
 Figure 945: DNA324799, NP\_061823.1, 204806.x.at  
 Figure 946: PRO81414  
 Figure 947: DNA154704, DNA154704, 204807.at  
 Figure 948: DNA328544, NP\_006673.1, 204834.at  
 Figure 949: PRO84347  
 Figure 950: DNA225661, NP\_001944.1, 204858.s.at

- Figure 951: PRO36124  
 Figure 952: DNA328545, NP\_064525.1, 204859.s.at  
 Figure 953: PRO84348  
 Figure 954A-B: DNA227629, NP\_004527.1, 204860.s.at  
 Figure 955: PRO38092  
 Figure 956: DNA328546, NP\_005249.1, 204867.at  
 Figure 957: PRO84349  
 Figure 958: DNA255993, NP\_008936.1, 204872.at  
 Figure 959: PRO51044  
 Figure 960: DNA273666, NP\_003349.1, 204881.s.at  
 Figure 961: PRO61634  
 Figure 962A-B: DNA76503, NP\_001549.1, 204912.at  
 Figure 963: PRO2536  
 Figure 964: DNA328547, TLR2, 204924.at  
 Figure 965: PRO208  
 Figure 966: DNA228014, NP\_002153.1, 204949.at  
 Figure 967: PRO38477  
 Figure 968: DNA328548, NP\_006298.1, 204955.at  
 Figure 969: PRO2618  
 Figure 970: DNA103283, NP\_002423.1, 204959.at  
 Figure 971: PRO4613  
 Figure 972: DNA227091, NP\_000256.1, 204961.s.at  
 Figure 973: PRO37554  
 Figure 974A-B: DNA328549, NP\_002897.1, 204969.s.at  
 Figure 975: PRO84350  
 Figure 976: DNA328301, NP\_005204.1, 204971.at  
 Figure 977: PRO70371  
 Figure 978A-B: DNA328550, NP\_001439.2, 204983.s.at  
 Figure 979: PRO937  
 Figure 980: DNA269665, NP\_002454.1, 204994.at  
 Figure 981: PRO58076  
 Figure 982A-B: DNA273686, NP\_055520.1, 205003.at  
 Figure 983: PRO61653  
 Figure 984: DNA272427, NP\_004799.1, 205005.s.at  
 Figure 985: PRO60679  
 Figure 986: DNA194830, NP\_055437.1, 205011.at  
 Figure 987: PRO24094  
 Figure 988: DNA328551, NP\_003823.1, 205048.s.at  
 Figure 989: PRO84351  
 Figure 990A-B: DNA328552, NP\_055886.1, 205068.s.at  
 Figure 991: PRO84352  
 Figure 992: DNA328553, NP\_061944.1, 205070.at  
 Figure 993: PRO84353  
 Figure 994: DNA194627, NP\_003051.1, 205074.at  
 Figure 995: PRO23962  
 Figure 996: DNA272181, NP\_006688.1, 205076.s.at  
 Figure 997: PRO60446  
 Figure 998: DNA254216, NP\_002020.1, 205119.s.at  
 Figure 999: PRO49328  
 Figure 1000: DNA299899, NP\_002148.1, 205133.s.at  
 Figure 1001: PRO62760  
 Figure 1002: DNA328554, NP\_038202.1, 205147.x.at  
 Figure 1003: PRO84354  
 Figure 1004: DNA328555, NP\_001241.1, 205153.s.at  
 Figure 1005: PRO34457  
 Figure 1006: DNA80896, NP\_001100.1, 205180.s.at  
 Figure 1007: PRO1686  
 Figure 1008: DNA328556, NP\_004568.1, 205194.at  
 Figure 1009: PRO84355  
 Figure 1010: DNA273535, NP\_004217.1, 205214.at  
 Figure 1011: PRO61515  
 Figure 1012: DNA93504, NP\_006009.1, 205220.at  
 Figure 1013: PRO4923  
 Figure 1014: DNA325255, NP\_001994.2, 205237.at  
 Figure 1015: PRO1910  
 Figure 1016: DNA327634, NP\_005129.1, 205241.at  
 Figure 1017: PRO83636  
 Figure 1018: DNA227081, NP\_000390.2, 205249.at  
 Figure 1019: PRO37544  
 Figure 1020: DNA328557, NP\_001098.1, 205260.s.at  
 Figure 1021: PRO84356  
 Figure 1022: DNA328558, BC016618, 205269.at  
 Figure 1023: PRO84357  
 Figure 1024: DNA328559, NP\_005556.1, 205270.s.at  
 Figure 1025: PRO84358  
 Figure 1026A-B: DNA227505, NP\_003670.1, 205306.x.at  
 Figure 1027: PRO37968  
 Figure 1028: DNA325783, NP\_002558.1, 205353.s.at  
 Figure 1029: PRO59001  
 Figure 1030: DNA88215, NP\_001919.1, 205382.s.at  
 Figure 1031: PRO2703  
 Figure 1032: DNA328560, NP\_003650.1, 205401.at  
 Figure 1033: PRO84359  
 Figure 1034: DNA328561, NP\_004624.1, 205403.at  
 Figure 1035: PRO2019  
 Figure 1036: DNA327638, NP\_005516.1, 205404.at  
 Figure 1037: PRO83639  
 Figure 1038: DNA328562, NP\_000010.1, 205412.at  
 Figure 1039: PRO84360  
 Figure 1040A-B: DNA328563, NP\_005329.2, 205425.at  
 Figure 1041: PRO81554  
 Figure 1042: DNA328564, HPCAL1, 205462.s.at  
 Figure 1043: PRO84361  
 Figure 1044: DNA196825, NP\_005105.1, 205466.s.at  
 Figure 1045: PRO25266  
 Figure 1046: DNA328565, NP\_057070.1, 205474.at  
 Figure 1047: PRO84362  
 Figure 1048: DNA226153, NP\_002649.1, 205479.s.at  
 Figure 1049: PRO36616  
 Figure 1050: DNA287224, NP\_005092.1, 205483.s.at  
 Figure 1051: PRO69503  
 Figure 1052: DNA328566, NP\_060446.1, 205510.s.at  
 Figure 1053: PRO84363  
 Figure 1054: DNA328567, NP\_006797.2, 205548.s.at  
 Figure 1055: PRO84364  
 Figure 1056: DNA227535, NP\_066190.1, 205568.at



- Figure 1057: PRO37998  
Figure 1058A-B: DNA327643, NP\_055712.1, 205594.at  
Figure 1059: PRO83644  
Figure 1060A-C: DNA328568, NP\_006720.1, 205603.s.at  
Figure 1061: PRO59731  
Figure 1062: DNA324324, NP\_000679.1, 205633.s.at  
Figure 1063: PRO81000  
Figure 1064: DNA328569, NP\_077274.1, 205634.x.at  
Figure 1065: PRO84365  
Figure 1066: DNA88076, NP\_001628.1, 205639.at  
Figure 1067: PRO2640  
Figure 1068: DNA287317, NP\_003724.1, 205660.at  
Figure 1069: PRO69582  
Figure 1070: DNA328570, NP\_004040.1, 205681.at  
Figure 1071: PRO37843  
Figure 1072: DNA327644, NP\_060395.2, 205684.s.at  
Figure 1073: PRO83645  
Figure 1074: DNA150621, NP\_036595.1, 205704.s.at  
Figure 1075: PRO12374  
Figure 1076: DNA328571, NP\_001254.1, 205709.s.at  
Figure 1077: PRO84366  
Figure 1078: DNA88106, NP\_004325.1, 205715.at  
Figure 1079: PRO2655  
Figure 1080: DNA270401, NP\_003140.1, 205743.at  
Figure 1081: PRO58784  
Figure 1082: DNA275620, NP\_000628.1, 205770.at  
Figure 1083: PRO63244  
Figure 1084: DNA88187, NP\_001757.1, 205789.at  
Figure 1085: PRO2689  
Figure 1086: DNA76517, NP\_002176.1, 205798.at  
Figure 1087: PRO2541  
Figure 1088A-B: DNA271915, NP\_056191.1, 205801.s.at  
Figure 1089: PRO60192  
Figure 1090: DNA194766, NP\_079504.1, 205804.s.at  
Figure 1091: PRO24046  
Figure 1092: DNA328572, NP\_004309.2, 205808.at  
Figure 1093: PRO84367  
Figure 1094: DNA328573, NP\_006761.1, 205819.at  
Figure 1095: PRO1559  
Figure 1096A-B: DNA328574, NP\_004963.1, 205842.s.at  
Figure 1097: PRO84368  
Figure 1098: DNA327651, NP\_005612.1, 205863.at  
Figure 1099: PRO83649  
Figure 1100: DNA328575, NP\_071754.2, 205872.x.at  
Figure 1101: PRO84369  
Figure 1102A-B: DNA220746, NP\_000876.1, 205884.at  
Figure 1103: PRO34724  
Figure 1104A-B: DNA273962, NP\_055605.1, 205888.s.at  
Figure 1105: PRO61910  
Figure 1106: DNA93423, NP\_000667.1, 205891.at  
Figure 1107: PRO4944  
Figure 1108: DNA328576, HSU20350, 205898.at  
Figure 1109: PRO4940  
Figure 1110: DNA328577, NP\_003905.1, 205899.at  
Figure 1111: PRO59588  
Figure 1112A-B: DNA196549, NP\_003034.1, 205920.at  
Figure 1113: PRO25031  
Figure 1114: DNA328578, NP\_004656.2, 205922.at  
Figure 1115: PRO7426  
Figure 1116A-B: DNA270867, NP\_006217.1, 205934.at  
Figure 1117: PRO59203  
Figure 1118: DNA76516, NP\_000556.1, 205945.at  
Figure 1119: PRO2022  
Figure 1120: DNA196439, NP\_003865.1, 205988.at  
Figure 1121: PRO24934  
Figure 1122: DNA36722, NP\_000576.1, 205992.s.at  
Figure 1123: PRO77  
Figure 1124: DNA328579, BC020082, 206020.at  
Figure 1125: PRO84370  
Figure 1126: DNA328580, HSU27699, 206058.at  
Figure 1127: PRO4627  
Figure 1128: DNA328581, NP\_002122.1, 206074.s.at  
Figure 1129: PRO34536  
Figure 1130: DNA328582, NP\_001865.1, 206100.at  
Figure 1131: PRO84371  
Figure 1132: DNA226105, NP\_002925.1, 206111.at  
Figure 1133: PRO36568  
Figure 1134: DNA225764, NP\_000037.1, 206129.s.at  
Figure 1135: PRO36227  
Figure 1136: DNA328583, ASGR2, 206130.s.at  
Figure 1137: PRO84372  
Figure 1138: DNA327656, NP\_055294.1, 206134.at  
Figure 1139: PRO36117  
Figure 1140A-B: DNA271837, NP\_055497.1, 206135.at  
Figure 1141: PRO60117  
Figure 1142: DNA328584, NP\_001148.1, 206200.s.at  
Figure 1143: PRO4833  
Figure 1144: DNA226058, NP\_005075.1, 206214.at  
Figure 1145: PRO36521  
Figure 1146: DNA218691, NP\_003832.1, 206222.at  
Figure 1147: PRO34469  
Figure 1148A-C: DNA328585, AF286028, 206239.s.at  
Figure 1149: DNA328586, NP\_002369.2, 206267.s.at  
Figure 1150: PRO84373  
Figure 1151: DNA328587, NP\_002612.1, 206380.s.at  
Figure 1152: PRO2854  
Figure 1153: DNA255814, NP\_005840.1, 206420.at  
Figure 1154: PRO50869  
Figure 1155: DNA328588, NP\_060823.1, 206500.s.at  
Figure 1156: PRO84374  
Figure 1157: DNA270444, NP\_004824.1, 206513.at  
Figure 1158: PRO58823

- Figure 1159: DNA196614, NP\_001158.1, 206536.s.at  
 Figure 1160: PRO25091  
 Figure 1161: DNA270019, NP\_036351.1, 206538.at  
 Figure 1162: PRO58414  
 Figure 1163: DNA327663, NP\_006771.1, 206565.x.at  
 Figure 1164: PRO83654  
 Figure 1165: DNA327665, NP\_002099.1, 206643.at  
 Figure 1166: PRO83655  
 Figure 1167: DNA328589, BCL2L1, 206665.s.at  
 Figure 1168: PRO83141  
 Figure 1169: DNA328590, C6orf32, 206707.x.at  
 Figure 1170: PRO84375  
 Figure 1171A-B: DNA88191, NP\_001234.1, 206729.at  
 Figure 1172: PRO2691  
 Figure 1173: DNA327669, NP\_000914.1, 206792.x.at  
 Figure 1174: PRO83657  
 Figure 1175: DNA270107, NP\_006856.1, 206881.s.at  
 Figure 1176: PRO58498  
 Figure 1177: DNA256561, NP\_062550.1, 206914.at  
 Figure 1178: PRO51592  
 Figure 1179: DNA328591, NP\_006635.1, 206976.s.at  
 Figure 1180: PRO84376  
 Figure 1181A-B: DNA227659, NP\_000570.1, 206991.s.at  
 Figure 1182: PRO38122  
 Figure 1183: DNA188289, NP\_001548.1, 207008.at  
 Figure 1184: PRO21820  
 Figure 1185: DNA328592, AB015228, 207016.s.at  
 Figure 1186: PRO84377  
 Figure 1187: DNA227531, NP\_004722.1, 207057.at  
 Figure 1188: PRO37994  
 Figure 1189: DNA327673, NP\_002188.1, 207071.s.at  
 Figure 1190: PRO83660  
 Figure 1191A-B: DNA328593, CIAS1, 207075.at  
 Figure 1192: PRO84378  
 Figure 1193A-B: DNA328594, CSF1, 207082.at  
 Figure 1194: PRO84379  
 Figure 1195: DNA88291, NP\_001965.1, 207111.at  
 Figure 1196: PRO2729  
 Figure 1197A-B: DNA327674, NP\_002739.1, 207121.s.at  
 Figure 1198: PRO83661  
 Figure 1199: DNA328595, NP\_001045.1, 207122.x.at  
 Figure 1200: PRO84380  
 Figure 1201: DNA226996, NP\_000239.1, 207233.s.at  
 Figure 1202: PRO37459  
 Figure 1203A-B: DNA226536, NP\_003225.1, 207332.s.at  
 Figure 1204: PRO36999  
 Figure 1205: DNA227668, NP\_000158.1, 207387.s.at  
 Figure 1206: PRO38131  
 Figure 1207: DNA328596, DEGS, 207431.s.at  
 Figure 1208: PRO37741  
 Figure 1209: DNA274829, NP\_003653.1, 207469.s.at  
 Figure 1210: PRO62588  
 Figure 1211: DNA328597, NP\_001680.1, 207507.s.at  
 Figure 1212: PRO84381  
 Figure 1213: DNA328598, NP\_055146.1, 207528.s.at  
 Figure 1214: PRO23276  
 Figure 1215: DNA328599, NFKB2, 207535.s.at  
 Figure 1216: PRO84382  
 Figure 1217: DNA328600, NP\_004839.1, 207571.x.at  
 Figure 1218: PRO84383  
 Figure 1219: DNA328601, NP\_056490.1, 207574.s.at  
 Figure 1220: PRO84384  
 Figure 1221: DNA328602, NP\_002261.1, 207657.x.at  
 Figure 1222: PRO84385  
 Figure 1223: DNA226278, NP\_005865.1, 207697.x.at  
 Figure 1224: PRO36741  
 Figure 1225: DNA227395, NP\_005331.1, 207721.x.at  
 Figure 1226: PRO37858  
 Figure 1227: DNA325654, NP\_054752.1, 207761.s.at  
 Figure 1228: PRO4348  
 Figure 1229: DNA226930, NP\_004152.1, 207791.s.at  
 Figure 1230: PRO37393  
 Figure 1231: DNA328603, NP\_000304.1, 207808.s.at  
 Figure 1232: PRO84386  
 Figure 1233: DNA328604, NP\_001174.2, 207809.s.at  
 Figure 1234: PRO84387  
 Figure 1235: DNA327682, NP\_001905.1, 207843.x.at  
 Figure 1236: PRO83666  
 Figure 1237: DNA36708, NP\_002081.1, 207850.at  
 Figure 1238: PRO34256  
 Figure 1239: DNA199788, NP\_002981.1, 207861.at  
 Figure 1240: PRO34107  
 Figure 1241: DNA328605, ST7, 207871.s.at  
 Figure 1242: PRO84388  
 Figure 1243: DNA256523, NP\_006854.1, 207872.s.at  
 Figure 1244: PRO51557  
 Figure 1245: DNA218651, NP\_003798.1, 207907.at  
 Figure 1246: PRO34447  
 Figure 1247: DNA275286, NP\_009205.1, 208002.s.at  
 Figure 1248: PRO62967  
 Figure 1249A-B: DNA328606, CBFA2T3, 208056.s.at  
 Figure 1250: PRO84389  
 Figure 1251A-B: DNA328607, NP\_003639.1, 208072.s.at  
 Figure 1252: PRO84390  
 Figure 1253: DNA327685, NP\_067586.1, 208074.s.at  
 Figure 1254: PRO83669  
 Figure 1255: DNA328608, NP\_006264.2, 208075.s.at  
 Figure 1256: PRO9932  
 Figure 1257: DNA255376, NP\_110423.1, 208091.s.at  
 Figure 1258: PRO50444  
 Figure 1259: DNA327686, NP\_005898.1, 208116.s.at  
 Figure 1260: PRO83670  
 Figure 1261A-B: DNA328609, NP\_109592.1, 208121.s.at  
 Figure 1262: PRO84391  
 Figure 1263: DNA328610, NP\_112601.1, 208146.s.at  
 Figure 1264: PRO84392  
 Figure 1265A-B: DNA226706, NP\_003777.2,

- 208161.s.at  
 Figure 1266: PRO37169  
 Figure 1267: DNA328611, RASGRP2, 208206.s.at  
 Figure 1268: PRO84393  
 Figure 1269: DNA328612, NP\_000166.2, 208308.s.at  
 Figure 1270: PRO84394  
 Figure 1271: DNA270558, NP\_006734.1, 208319.s.at  
 Figure 1272: PRO58933  
 Figure 1273: DNA227614, NP\_004859.1, 208336.s.at  
 Figure 1274: PRO38077  
 Figure 1275: DNA327690, NP\_004022.1, 208436.s.at  
 Figure 1276: PRO83673  
 Figure 1277: DNA328613, NP\_056953.2, 208510.s.at  
 Figure 1278: PRO84395  
 Figure 1279A-C: DNA328614, SRRM2, 208610.s.at  
 Figure 1280: PRO84396  
 Figure 1281A-C: DNA328615, NP\_003118.1, 208611.s.at  
 Figure 1282: PRO84397  
 Figure 1283A-C: DNA328616, NP\_001448.1, 208613.s.at  
 Figure 1284: PRO84398  
 Figure 1285: DNA326362, VATI, 208626.s.at  
 Figure 1286: PRO82758  
 Figure 1287: DNA325912, NP\_001093.1, 208637.x.at  
 Figure 1288: PRO82367  
 Figure 1289: DNA271268, NP\_009057.1, 208649.s.at  
 Figure 1290: PRO59579  
 Figure 1291: DNA328617, AF299343, 208653.s.at  
 Figure 1292: PRO84399  
 Figure 1293A-C: DNA328618, NP\_003307.2, 208664.s.at  
 Figure 1294: PRO84400  
 Figure 1295: DNA304686, NP\_002565.1, 208680.at  
 Figure 1296: PRO71112  
 Figure 1297: DNA304499, NP\_006588.1, 208687.x.at  
 Figure 1298: PRO71063  
 Figure 1299A-B: DNA328619, BC001188, 208691.at  
 Figure 1300: PRO84401  
 Figure 1301: DNA287189, NP\_002038.1, 208693.s.at  
 Figure 1302: PRO69475  
 Figure 1303: DNA324217, ATIC, 208758.at  
 Figure 1304: PRO80908  
 Figure 1305: DNA327696, AF228339, 208763.s.at  
 Figure 1306: PRO83679  
 Figure 1307: DNA328620, AK000295, 208772.at  
 Figure 1308: PRO84402  
 Figure 1309: DNA328621, NP\_002788.1, 208799.at  
 Figure 1310: PRO84403  
 Figure 1311: DNA287169, CAA42052.1, 208805.at  
 Figure 1312: PRO10404  
 Figure 1313: DNA324531, NP\_002120.1, 208808.s.at  
 Figure 1314: PRO81185  
 Figure 1315: DNA273521, NP\_002070.1, 208813.at  
 Figure 1316: PRO61502  
 Figure 1317: DNA328622, BC000835, 208827.at  
 Figure 1318: PRO82662  
 Figure 1319: DNA227556, NP\_001670.1, 208836.at  
 Figure 1320: PRO38019  
 Figure 1321: DNA326042, NP\_031390.1, 208837.at  
 Figure 1322: PRO1078  
 Figure 1323A-B: DNA328623, NP\_056107.1, 208858.s.at  
 Figure 1324: PRO61321  
 Figure 1325: DNA227874, NP\_003320.1, 208864.s.at  
 Figure 1326: PRO38337  
 Figure 1327: DNA328624, BC003562, 208891.at  
 Figure 1328: PRO59076  
 Figure 1329: DNA328625, NP\_073143.1, 208892.s.at  
 Figure 1330: PRO84404  
 Figure 1331: DNA328626, NP\_057078.1, 208898.at  
 Figure 1332: PRO61768  
 Figure 1333: DNA327700, BC015130, 208905.at  
 Figure 1334: PRO83683  
 Figure 1335: DNA325472, NP\_116056.2, 208906.at  
 Figure 1336: PRO81995  
 Figure 1337A-B: DNA328627, FLJ13052, 208918.s.at  
 Figure 1338: PRO84405  
 Figure 1339: DNA325473, NP\_006353.2, 208922.s.at  
 Figure 1340: PRO81996  
 Figure 1341: DNA287238, NP\_000425.1, 208926.at  
 Figure 1342: PRO69515  
 Figure 1343: DNA328628, NP\_060542.2, 208933.s.at  
 Figure 1344: PRO84406  
 Figure 1345: DNA290261, NP\_001291.2, 208960.s.at  
 Figure 1346: PRO70387  
 Figure 1347A-B: DNA325478, NP\_037534.2, 208962.s.at  
 Figure 1348: PRO81999  
 Figure 1349: DNA328629, NP\_006079.1, 208977.x.at  
 Figure 1350: PRO84407  
 Figure 1351: DNA328630, NP\_036293.1, 209004.s.at  
 Figure 1352: PRO84408  
 Figure 1353: DNA328631, AK027318, 209006.s.at  
 Figure 1354: PRO84409  
 Figure 1355: DNA328632, DJ465N24.2.1Homo, 209007.s.at  
 Figure 1356: DNA328633, NP\_004784.2, 209017.s.at  
 Figure 1357: PRO84411  
 Figure 1358A-B: DNA328634, NP\_006594.1, 209023.s.at  
 Figure 1359: PRO84412  
 Figure 1360: DNA328635, BC020946, 209026.x.at  
 Figure 1361: PRO84413  
 Figure 1362: DNA274202, NP\_006804.1, 209034.at  
 Figure 1363: PRO62131  
 Figure 1364: DNA328636, PAPSS1, 209043.at  
 Figure 1365: PRO84414  
 Figure 1366A-C: DNA328637, HSA7042, 209053.s.at  
 Figure 1367: PRO81109  
 Figure 1368: DNA326406, NP\_005315.1, 209069.s.at  
 Figure 1369: PRO11403

- Figure 1370: DNA227289, NP\_006532.1, 209080.x.at  
 Figure 1371: PRO37752  
 Figure 1372: DNA274180, NP\_009005.1, 209083.at  
 Figure 1373: PRO62110  
 Figure 1374: DNA327707, NP\_000148.1, 209093.s.at  
 Figure 1375: PRO83689  
 Figure 1376: DNA226564, NP\_000099.1, 209095.at  
 Figure 1377: PRO37027  
 Figure 1378: DNA325163, NP\_001113.1, 209122.at  
 Figure 1379: PRO81730  
 Figure 1380: DNA328638, BC000576, 209123.at  
 Figure 1381: PRO81129  
 Figure 1382: DNA274723, AAB62222.1, 209129.at  
 Figure 1383: PRO62502  
 Figure 1384: DNA328639, HSM801840, 209132.s.at  
 Figure 1385: PRO84415  
 Figure 1386: DNA328640, ASPH, 209135.at  
 Figure 1387: PRO84416  
 Figure 1388: DNA327713, BC010653, 209146.at  
 Figure 1389: PRO37975  
 Figure 1390: DNA271937, NP\_055419.1, 209154.at  
 Figure 1391: PRO60213  
 Figure 1392: DNA328641, NP\_001840.2, 209156.s.at  
 Figure 1393: PRO84417  
 Figure 1394: DNA325285, AKR1C3, 209160.at  
 Figure 1395: PRO81832  
 Figure 1396A-B: DNA328642, AF073310, 209184.s.at  
 Figure 1397: PRO84418  
 Figure 1398A-B: DNA328643, HUMHK1A, 209186.at  
 Figure 1399: PRO84419  
 Figure 1400: DNA189700, NP\_005243.1, 209189.at  
 Figure 1401: PRO25619  
 Figure 1402: DNA327715, NP\_115914.1, 209191.at  
 Figure 1403: PRO83694  
 Figure 1404: DNA103520, NP\_002639.1, 209193.at  
 Figure 1405: PRO4847  
 Figure 1406A-B: DNA269816, MEF2C, 209199.s.at  
 Figure 1407: PRO58219  
 Figure 1408: DNA328644, 349746.9, 209200.at  
 Figure 1409: PRO84420  
 Figure 1410: DNA326891, NP\_001748.1, 209213.at  
 Figure 1411: PRO83212  
 Figure 1412: DNA328645, NP\_009006.1, 209216.at  
 Figure 1413: PRO84421  
 Figure 1414: DNA227483, NP\_003120.1, 209218.at  
 Figure 1415: PRO37946  
 Figure 1416: DNA328646, NP\_036517.1, 209230.s.at  
 Figure 1417: PRO84422  
 Figure 1418A-C: DNA328647, AB017133, 209234.at  
 Figure 1419: PRO84423  
 Figure 1420A-B: DNA328648, D87075, 209236.at  
 Figure 1421: DNA328649, NP\_116093.1, 209251.x.at  
 Figure 1422: PRO84424  
 Figure 1423: DNA255255, NP\_071437.1, 209267.s.at  
 Figure 1424: PRO50332  
 Figure 1425A-B: DNA226827, NP\_001673.1, 209281.s.at  
 Figure 1426: PRO37290  
 Figure 1427: DNA328650, 200118.10, 209286.at  
 Figure 1428: PRO84425  
 Figure 1429: DNA274883, NP\_000058.1, 209301.at  
 Figure 1430: PRO62628  
 Figure 1431: DNA328651, AF087853, 209305.s.at  
 Figure 1432: PRO82889  
 Figure 1433: DNA327718, CASP4, 209310.s.at  
 Figure 1434: PRO83697  
 Figure 1435: DNA328652, NP\_077298.1, 209321.s.at  
 Figure 1436: PRO84426  
 Figure 1437: DNA328653, AF063020, 209337.at  
 Figure 1438: PRO84427  
 Figure 1439: DNA328654, UAP1, 209340.at  
 Figure 1440: PRO84428  
 Figure 1441: DNA328655, 346677.3, 209341.s.at  
 Figure 1442: PRO84429  
 Figure 1443: DNA269630, NP\_003281.1, 209344.at  
 Figure 1444: PRO58042  
 Figure 1445A-B: DNA328656, HSA303098, 209345.s.at  
 Figure 1446: PRO84430  
 Figure 1447A-B: DNA328657, NP\_060895.1, 209346.s.at  
 Figure 1448: PRO84431  
 Figure 1449A-B: DNA328658, AF055376, 209348.s.at  
 Figure 1450: PRO84432  
 Figure 1451: DNA327719, NP\_003704.2, 209355.s.at  
 Figure 1452: PRO83698  
 Figure 1453: DNA328659, ECM1, 209365.s.at  
 Figure 1454: PRO84433  
 Figure 1455: DNA225952, NP\_001267.1, 209395.at  
 Figure 1456: PRO36415  
 Figure 1457: DNA275366, BC001851, 209444.at  
 Figure 1458: PRO63036  
 Figure 1459: DNA328660, NP\_003675.2, 209467.s.at  
 Figure 1460: PRO84434  
 Figure 1461A-B: DNA328661, NP\_006304.1, 209475.at  
 Figure 1462: PRO84435  
 Figure 1463: DNA328662, OSBPL1A, 209485.s.at  
 Figure 1464: PRO84436  
 Figure 1465: DNA324899, NP\_002938.1, 209507.at  
 Figure 1466: PRO81503  
 Figure 1467: DNA274027, HSU38654, 209515.s.at  
 Figure 1468: PRO61971  
 Figure 1469: DNA328663, NP\_057157.1, 209524.at  
 Figure 1470: PRO36183  
 Figure 1471A-C: DNA328664, NP\_009131.1, 209534.x.at  
 Figure 1472: PRO84437  
 Figure 1473A-B: DNA328665, RGL, 209568.s.at

- Figure 1474: PRO84438  
 Figure 1475: DNA328666, AF084943, 209585.s.at  
 Figure 1476: PRO1917  
 Figure 1477: DNA328667, S69189, 209600.s.at  
 Figure 1478: PRO84439  
 Figure 1479: DNA328668, NP\_003157.1, 209607.x.at  
 Figure 1480: PRO84440  
 Figure 1481: DNA328669, NP\_005882.1, 209608.s.at  
 Figure 1482: PRO84441  
 Figure 1483A-B: DNA328670, BC001618, 209610.s.at  
 Figure 1484: PRO70011  
 Figure 1485: DNA256209, NP\_002259.1, 209653.at  
 Figure 1486: PRO51256  
 Figure 1487A-B: DNA272671, HSU26710, 209682.at  
 Figure 1488: PRO60796  
 Figure 1489: DNA151564, DNA151564, 209683.at  
 Figure 1490: PRO11886  
 Figure 1491: DNA327727, NP\_000308.1, 209694.at  
 Figure 1492: PRO83705  
 Figure 1493: DNA328671, NP\_000498.2, 209696.at  
 Figure 1494: PRO84442  
 Figure 1495: DNA327728, BC004492, 209703.x.at  
 Figure 1496: PRO4348  
 Figure 1497: DNA328672, CAA68871.1, 209707.at  
 Figure 1498: PRO84444  
 Figure 1499A-B: DNA328673, HUMCSDF1, 209716.at  
 Figure 1500: PRO84445  
 Figure 1501A-B: DNA304800, BC002538, 209723.at  
 Figure 1502: PRO69458  
 Figure 1503A-B: DNA328674, NP\_056011.1, 209760.at  
 Figure 1504: PRO84446  
 Figure 1505: DNA324250, NP\_536349.1, 209761.s.at  
 Figure 1506: PRO80934  
 Figure 1507A-B: DNA328675, ADAM19, 209765.at  
 Figure 1508: PRO84447  
 Figure 1509: DNA327731, NP\_003302.1, 209803.s.at  
 Figure 1510: PRO83707  
 Figure 1511: DNA328676, IL16, 209827.s.at  
 Figure 1512: PRO84448  
 Figure 1513A-B: DNA196499, AB002384, 209829.at  
 Figure 1514: PRO24988  
 Figure 1515: DNA328677, AF060511, 209836.x.at  
 Figure 1516: PRO84449  
 Figure 1517: DNA324805, NP\_008978.1, 209846.s.at  
 Figure 1518: PRO81419  
 Figure 1519: DNA273915, NP\_036215.1, 209864.at  
 Figure 1520: PRO61867  
 Figure 1521: DNA290585, NP\_000573.1, 209875.s.at  
 Figure 1522: PRO70536  
 Figure 1523: DNA328678, NP\_008843.1, 209882.at  
 Figure 1524: PRO62586  
 Figure 1525: DNA328679, 347423.1, 209892.at  
 Figure 1526: PRO84450  
 Figure 1527: DNA328258, HSM802616, 209900.s.at  
 Figure 1528: PRO84151  
 Figure 1529A-B: DNA328680, NP\_062541.1, 209907.s.at  
 Figure 1530: PRO84451  
 Figure 1531: DNA299884, AB040875, 209921.at  
 Figure 1532: PRO70858  
 Figure 1533: DNA328681, NP\_005089.1, 209928.s.at  
 Figure 1534: PRO84452  
 Figure 1535: DNA272326, NP\_006154.1, 209930.s.at  
 Figure 1536: PRO60583  
 Figure 1537: DNA328682, AF225981, 209935.at  
 Figure 1538: PRO84453  
 Figure 1539: DNA327754, NP\_150634.1, 209970.x.at  
 Figure 1540: PRO4526  
 Figure 1541: DNA328683, NP\_000399.1, 210007.s.at  
 Figure 1542: PRO84454  
 Figure 1543: DNA227660, NP\_001327.1, 210042.s.at  
 Figure 1544: PRO38123  
 Figure 1545: DNA327739, AF092535, 210058.at  
 Figure 1546: PRO83714  
 Figure 1547: DNA327740, NP\_003944.1, 210087.s.at  
 Figure 1548: PRO1787  
 Figure 1549: DNA328684, BC001234, 210102.at  
 Figure 1550: PRO84455  
 Figure 1551A-B: DNA328685, NP\_127497.1, 210113.s.at  
 Figure 1552: PRO34751  
 Figure 1553: DNA328686, NP\_000566.1, 210118.s.at  
 Figure 1554: PRO64  
 Figure 1555: DNA227757, NP\_000743.1, 210128.s.at  
 Figure 1556: PRO38220  
 Figure 1557: DNA227501, NP\_000295.1, 210139.s.at  
 Figure 1558: PRO37964  
 Figure 1559: DNA328687, AF004231, 210146.x.at  
 Figure 1560: PRO84456  
 Figure 1561A-B: DNA328688, NP\_006838.2, 210152.at  
 Figure 1562: PRO84457  
 Figure 1563: DNA328689, NP\_003259.2, 210166.at  
 Figure 1564: PRO7521  
 Figure 1565: DNA270196, HUMZFM1B, 210172.at  
 Figure 1566: PRO58584  
 Figure 1567: DNA328690, NP\_524145.1, 210240.s.at  
 Figure 1568: PRO59660  
 Figure 1569: DNA326963, HRIHFB2122, 210276.s.at  
 Figure 1570: PRO83276  
 Figure 1571: DNA328691, NP\_065717.1, 210346.s.at  
 Figure 1572: PRO84458  
 Figure 1573: DNA227652, NP\_002549.1, 210401.at  
 Figure 1574: PRO38115  
 Figure 1575: DNA225514, NP\_003864.1, 210510.s.at  
 Figure 1576: PRO35977  
 Figure 1577: DNA216517, NP\_005055.1, 210549.s.at  
 Figure 1578: PRO34269  
 Figure 1579: DNA327746, HUMGCBA, 210589.s.at

- Figure 1580: PRO83720  
 Figure 1581: DNA328692, AF025529, 210660.at  
 Figure 1582: PRO84459  
 Figure 1583: DNA272127, NP\_003928.1, 210663.s.at  
 Figure 1584: PRO60397  
 Figure 1585: DNA326525, NP\_006330.1, 210719.s.at  
 Figure 1586: PRO82894  
 Figure 1587: DNA226183, NP\_001453.1, 210773.s.at  
 Figure 1588: PRO36646  
 Figure 1589: DNA226078, NP\_000296.1, 210830.s.at  
 Figure 1590: PRO36541  
 Figure 1591: DNA226152, NP\_002650.1, 210845.s.at  
 Figure 1592: PRO36615  
 Figure 1593: DNA328693, HSU03891, 210873.x.at  
 Figure 1594: PRO84460  
 Figure 1595: DNA328694, BC007810, 210944.s.at  
 Figure 1596: PRO84461  
 Figure 1597: DNA213676, NP\_004604.1, 211003.x.at  
 Figure 1598: PRO35142  
 Figure 1599: DNA328695, NP\_002145.1, 211015.s.at  
 Figure 1600: PRO61480  
 Figure 1601: DNA328696, NP\_009214.1, 211026.s.at  
 Figure 1602: PRO62720  
 Figure 1603: DNA328697, NP\_116112.1, 211038.s.at  
 Figure 1604: PRO84462  
 Figure 1605: DNA328698, BC006403, 211063.s.at  
 Figure 1606: PRO12168  
 Figure 1607: DNA326712, NP\_001285.1, 211136.s.at  
 Figure 1608: PRO83054  
 Figure 1609A-B: DNA328699, AF189723, 211137.s.at  
 Figure 1610: PRO84463  
 Figure 1611: DNA327752, HSDHACTYL, 211150.s.at  
 Figure 1612A-B: DNA328700, SCD, 211162.x.at  
 Figure 1613: PRO84464  
 Figure 1614: DNA328701, PSEN2, 211373.s.at  
 Figure 1615: PRO80745  
 Figure 1616: DNA328702, NP\_036519.1, 211413.s.at  
 Figure 1617: PRO84465  
 Figure 1618: DNA256637, NP\_008849.1, 211423.s.at  
 Figure 1619: PRO51621  
 Figure 1620: DNA328703, NP\_003956.1, 211434.s.at  
 Figure 1621: PRO1873  
 Figure 1622: DNA327755, NP\_115957.1, 211458.s.at  
 Figure 1623: PRO83725  
 Figure 1624A-B: DNA328704, FGFR1, 211535.s.at  
 Figure 1625: PRO34231  
 Figure 1626: DNA324626, RIL, 211564.s.at  
 Figure 1627: PRO81272  
 Figure 1628: DNA328705, NP\_001345.1, 211653.x.at  
 Figure 1629: PRO62617  
 Figure 1630: DNA328706, BC021909, 211714.x.at  
 Figure 1631: PRO10347  
 Figure 1632A-B: DNA328707, AF172264, 211828.s.at  
 Figure 1633: PRO84466  
 Figure 1634: DNA226582, NP\_003863.1, 211844.s.at  
 Figure 1635: PRO37045  
 Figure 1636: DNA151912, BAA06683.1, 211935.at  
 Figure 1637: PRO12756  
 Figure 1638: DNA325941, NP\_005339.1, 211968.s.at  
 Figure 1639: PRO82388  
 Figure 1640: DNA287433, NP\_006810.1, 212009.s.at  
 Figure 1641: PRO69690  
 Figure 1642: DNA328708, NP\_002678.1, 212036.s.at  
 Figure 1643: PRO84467  
 Figure 1644: DNA103380, NP\_003365.1, 212038.s.at  
 Figure 1645: PRO4710  
 Figure 1646: DNA328709, BC004151, 212048.s.at  
 Figure 1647: PRO37676  
 Figure 1648A-B: DNA254751, AB018353, 212074.at  
 Figure 1649: DNA328710, HUMLAMA, 212086.x.at  
 Figure 1650A-B: DNA298616, NP\_001839.1, 212091.s.at  
 Figure 1651: PRO71027  
 Figure 1652: DNA154139, DNA154139, 212099.at  
 Figure 1653: DNA328711, AK023154, 212115.at  
 Figure 1654: PRO84468  
 Figure 1655: DNA328712, NP\_006501.1, 212118.at  
 Figure 1656: PRO84469  
 Figure 1657: DNA328713, AF100737, 212130.x.at  
 Figure 1658: PRO84470  
 Figure 1659: DNA328714, HSM801966, 212146.at  
 Figure 1660A-B: DNA151915, BAA09764.1, 212149.at  
 Figure 1661: PRO12758  
 Figure 1662: DNA88630, AAA52701.1, 212154.at  
 Figure 1663: PRO2877  
 Figure 1664: DNA328715, BC000950, 212160.at  
 Figure 1665: DNA328716, HSM800707, 212179.at  
 Figure 1666A-C: DNA255018, CAB61363.1, 212207.at  
 Figure 1667: PRO50107  
 Figure 1668A-B: DNA328717, CAB70761.1, 212232.at  
 Figure 1669: PRO84473  
 Figure 1670: DNA196116, DNA196116, 212246.at  
 Figure 1671A-B: DNA254262, NP\_055197.1, 212255.s.at  
 Figure 1672: PRO49373  
 Figure 1673: DNA327771, NP\_109591.1, 212268.at  
 Figure 1674: PRO83737  
 Figure 1675A-B: DNA328718, AAC39776.1, 212285.s.at  
 Figure 1676: PRO84474  
 Figure 1677: DNA328719, BC012895, 212295.s.at  
 Figure 1678: PRO84475  
 Figure 1679: DNA271103, NP\_005796.1, 212296.at  
 Figure 1680: PRO59425  
 Figure 1681A-B: DNA328720, HSA306929, 212297.at

Figure 1682: PRO84476  
 Figure 1683A-B: DNA328721, 1450005.12, 212298.at  
 Figure 1684: PRO84477  
 Figure 1685A-B: DNA150464, BAA34466.1, 212311.at  
 Figure 1686: PRO12270  
 Figure 1687: DNA326808, BC019307, 212312.at  
 Figure 1688: PRO83141  
 Figure 1689A-B: DNA124122, NP.005602.2, 212332.at  
 Figure 1690: PRO6323  
 Figure 1691: DNA287190, CAB43217.1, 212333.at  
 Figure 1692: PRO69476  
 Figure 1693A-B: DNA255527, HUMTI227HC, 212337.at  
 Figure 1694: DNA328722, BC012469, 212341.at  
 Figure 1695: PRO84478  
 Figure 1696: DNA328723, S47833, 212360.at  
 Figure 1697: PRO36682  
 Figure 1698A-B: DNA328724, AB007856, 212367.at  
 Figure 1699A-B: DNA327773, BAA25456.1, 212368.at  
 Figure 1700: PRO83739  
 Figure 1701A-C: DNA328725, AB007923, 212390.at  
 Figure 1702A-B: DNA150950, BAA07645.1, 212396.s.at  
 Figure 1703: PRO12554  
 Figure 1704A-B: DNA328726, BAA25466.2, 212443.at  
 Figure 1705: PRO84480  
 Figure 1706: DNA328727, AB033105, 212453.at  
 Figure 1707A-B: DNA328728, 481567.2, 212458.at  
 Figure 1708: PRO84482  
 Figure 1709: DNA151348, DNA151348, 212463.at  
 Figure 1710: PRO11726  
 Figure 1711A-: DNA328729, D80001, 212486.s.at  
 Figure 1712: PRO38526  
 Figure 1713A-B: DNA328730, BAA74899.2, 212492.s.at  
 Figure 1714: PRO84483  
 Figure 1715A-B: DNA328731, 234169.5, 212500.at  
 Figure 1716: PRO84484  
 Figure 1717: DNA328732, NP.116193.1, 212502.at  
 Figure 1718: PRO84485  
 Figure 1719: DNA0, AF038183, 212527.at  
 Figure 1720: PRO  
 Figure 1721: DNA328734, AAH01171.1, 212539.at  
 Figure 1722: PRO84487  
 Figure 1723: DNA328735, PHIP, 212542.s.at  
 Figure 1724: PRO84488  
 Figure 1725: DNA328736, BC009846, 212552.at  
 Figure 1726: PRO84489  
 Figure 1727A-D: DNA328737, 148650.1, 212560.at  
 Figure 1728: PRO84490  
 Figure 1729: DNA270260, HSPDCE2, 212568.s.at  
 Figure 1730A-B: DNA328738, BAA31625.1, 212569.at  
 Figure 1731: PRO84491  
 Figure 1732A-B: DNA328739, PTPRC, 212587.s.at  
 Figure 1733: PRO84492  
 Figure 1734: DNA327776, 1379302.1, 212593.s.at  
 Figure 1735: PRO83742  
 Figure 1736: DNA151487, DNA151487, 212594.at  
 Figure 1737: PRO11833  
 Figure 1738A-B: DNA328740, BAA76781.1, 212611.at  
 Figure 1739: PRO84493  
 Figure 1740: DNA81753, DNA81753, 212613.at  
 Figure 1741: PRO9216  
 Figure 1742A-B: DNA253817, BAA20767.1, 212615.at  
 Figure 1743: PRO49220  
 Figure 1744A-B: DNA328741, 474863.12, 212622.at  
 Figure 1745: PRO84494  
 Figure 1746: DNA194679, BAA05062.1, 212623.at  
 Figure 1747: PRO23989  
 Figure 1748A-B: DNA328742, 244522.6, 212628.at  
 Figure 1749: PRO59047  
 Figure 1750: DNA270683, NP.006247.1, 212629.s.at  
 Figure 1751: PRO59047  
 Figure 1752A-D: DNA327777, HSILIRECA, 212657.s.at  
 Figure 1753A-B: DNA150762, BAA13197.1, 212658.at  
 Figure 1754: PRO12455  
 Figure 1755: DNA327838, NP.000568.1, 212659.s.at  
 Figure 1756: PRO83789  
 Figure 1757: DNA328743, 1234685.2, 212667.at  
 Figure 1758: PRO84495  
 Figure 1759: DNA328744, AF318364, 212680.x.at  
 Figure 1760: PRO84496  
 Figure 1761: DNA328745, 482138.6, 212687.at  
 Figure 1762: PRO84497  
 Figure 1763: DNA324378, NP.000523.1, 212694.s.at  
 Figure 1764: PRO81047  
 Figure 1765: DNA328746, CAB43213.1, 212698.s.at  
 Figure 1766: PRO84498  
 Figure 1767A-B: DNA328747, BAA83030.1, 212765.at  
 Figure 1768: PRO84499  
 Figure 1769A-B: DNA328748, HSJ001388, 212774.at  
 Figure 1770: PRO59570  
 Figure 1771: DNA328749, HSM802266, 212779.at  
 Figure 1772: DNA328750, 7689361.1, 212812.at  
 Figure 1773: PRO84500  
 Figure 1774A-B: DNA328751, AF012086, 212842.x.at  
 Figure 1775: DNA328752, CAA76270.1, 212864.at  
 Figure 1776: PRO84501  
 Figure 1777A-B: DNA328753, BAA13212.1, 212873.at  
 Figure 1778: PRO84502

- Figure 1779: DNA271630, DNA271630, 212907.at  
 Figure 1780: DNA328754, 1397726.9, 212912.at  
 Figure 1781: PRO84503  
 Figure 1782A-B: DNA328755, BAA25490.1, 212946.at  
 Figure 1783: PRO84504  
 Figure 1784A-B: DNA328756, BAA74893.2, 212975.at  
 Figure 1785: PRO84505  
 Figure 1786: DNA154982, DNA154982, 213034.at  
 Figure 1787: DNA327785, BC017336, 213061.s.at  
 Figure 1788: PRO83749  
 Figure 1789A-C: DNA328757, 475076.9, 213069.at  
 Figure 1790: PRO84506  
 Figure 1791A-B: DNA328758, AB011123, 213109.at  
 Figure 1792: DNA272600, NP\_057259.1, 213112.s.at  
 Figure 1793: PRO60737  
 Figure 1794: DNA326217, NP\_004474.1, 213129.s.at  
 Figure 1795: PRO82630  
 Figure 1796: DNA228053, DNA228053, 213158.at  
 Figure 1797A-G: DNA103535, AF027153, 213164.at  
 Figure 1798: PRO4862  
 Figure 1799: DNA150875, CAB45717.1, 213246.at  
 Figure 1800: PRO11589  
 Figure 1801: DNA328759, HUMLPACI09, 213258.at  
 Figure 1802: DNA328760, 1376674.1, 213274.s.at  
 Figure 1803: PRO84508  
 Figure 1804A-B: DNA328761, BAA82991.1, 213280.at  
 Figure 1805: PRO84509  
 Figure 1806: DNA260974, NP\_006065.1, 213293.s.at  
 Figure 1807: PRO54720  
 Figure 1808: DNA328762, AAL30845.1, 213338.at  
 Figure 1809: PRO84510  
 Figure 1810: DNA327789, 1449824.5, 213348.at  
 Figure 1811: PRO83753  
 Figure 1812: DNA328763, NP\_001219.2, 213373.s.at  
 Figure 1813: PRO84511  
 Figure 1814: DNA328764, NP\_438169.1, 213375.s.at  
 Figure 1815: PRO84512  
 Figure 1816: DNA328765, 411350.1, 213391.at  
 Figure 1817: PRO84513  
 Figure 1818: DNA106195, DNA106195, 213454.at  
 Figure 1819: DNA327795, BC014226, 213457.at  
 Figure 1820: DNA328766, NP\_006077.1, 213476.x.at  
 Figure 1821: PRO84514  
 Figure 1822: DNA328767, BC008767, 213501.at  
 Figure 1823: PRO84515  
 Figure 1824: DNA254264, HSM800224, 213546.at  
 Figure 1825: PRO49375  
 Figure 1826: DNA328768, 1194561.1, 213572.s.at  
 Figure 1827: PRO84516  
 Figure 1828: DNA327800, 1251176.10, 213593.s.at  
 Figure 1829: PRO83763  
 Figure 1830: DNA151422, DNA151422, 213605.s.at  
 Figure 1831: PRO11792  
 Figure 1832: DNA225974, NP\_000864.1, 213620.s.at  
 Figure 1833: PRO36437  
 Figure 1834: DNA328769, CAA69330.1, 213624.at  
 Figure 1835: PRO84517  
 Figure 1836: DNA260173, DNA260173, 213638.at  
 Figure 1837: PRO54102  
 Figure 1838A-C: DNA273792, DNA273792, 213649.at  
 Figure 1839: DNA151886, CAB43234.1, 213682.at  
 Figure 1840: PRO12745  
 Figure 1841: DNA227788, NP\_002995.1, 213716.s.at  
 Figure 1842: PRO38251  
 Figure 1843: DNA328771, HSMYOSIE, 213733.at  
 Figure 1844: DNA328772, AAC19149.1, 213761.at  
 Figure 1845: PRO84519  
 Figure 1846: DNA328773, BC001528, 213766.x.at  
 Figure 1847: PRO84520  
 Figure 1848: DNA328774, NP\_004263.1, 213793.s.at  
 Figure 1849: PRO60536  
 Figure 1850A-B: DNA328775, NP\_006540.2, 213812.s.at  
 Figure 1851: PRO84521  
 Figure 1852: DNA328776, 407661.4, 213817.at  
 Figure 1853: PRO84522  
 Figure 1854A-B: DNA328777, IDN3, 213918.s.at  
 Figure 1855: PRO84523  
 Figure 1856: DNA196110, DNA196110, 214016.s.at  
 Figure 1857: PRO24635  
 Figure 1858: DNA150990, NP\_003632.1, 214022.s.at  
 Figure 1859: PRO12570  
 Figure 1860: DNA328778, 234498.37, 214093.s.at  
 Figure 1861: PRO84524  
 Figure 1862A-B: DNA272292, NP\_055459.1, 214130.s.at  
 Figure 1863: PRO60550  
 Figure 1864: DNA82378, NP\_002695.1, 214146.s.at  
 Figure 1865: PRO1725  
 Figure 1866A-B: DNA328779, 332730.12, 214155.s.at  
 Figure 1867: PRO84525  
 Figure 1868: DNA304659, NP\_002023.1, 214211.at  
 Figure 1869: PRO71086  
 Figure 1870: DNA256662, NP\_009112.1, 214219.x.at  
 Figure 1871: PRO51628  
 Figure 1872A-B: DNA328780, 480940.15, 214285.at  
 Figure 1873: PRO84526  
 Figure 1874: DNA328781, 1453703.13, 214349.at  
 Figure 1875: PRO84527  
 Figure 1876: DNA273174, NP\_001951.1, 214394.x.at  
 Figure 1877: PRO61211  
 Figure 1878: DNA328782, 337794.1, 214405.at  
 Figure 1879: PRO84528  
 Figure 1880: DNA287630, NP\_000160.1, 214430.at  
 Figure 1881: PRO2154  
 Figure 1882: DNA227376, NP\_005393.1, 214435.x.at  
 Figure 1883: PRO37839



- Figure 1884: DNA273138, NP\_005495.1, 214452.at  
 Figure 1885: PRO61182  
 Figure 1886: DNA327812, NP\_006408.2, 214453.s.at  
 Figure 1887: PRO83773  
 Figure 1888: DNA302598, NP\_066361.1, 214487.s.at  
 Figure 1889: PRO62511  
 Figure 1890: DNA328783, NP\_002021.2, 214560.at  
 Figure 1891: PRO84529  
 Figure 1892: DNA324728, BC017730, 214581.x.at  
 Figure 1893: PRO868  
 Figure 1894A-B: DNA328784, 331045.1, 214582.at  
 Figure 1895: PRO84530  
 Figure 1896: DNA328785, NP\_004062.1, 214683.s.at  
 Figure 1897: PRO84531  
 Figure 1898: DNA328786, BC017407, 214686.at  
 Figure 1899: PRO84532  
 Figure 1900: DNA271990, DNA271990, 214722.at  
 Figure 1901A-B: DNA274485, AB007863, 214735.at  
 Figure 1902: DNA328787, 238292.8, 214746.s.at  
 Figure 1903: PRO84533  
 Figure 1904: DNA328788, AK023937, 214763.at  
 Figure 1905: PRO29183  
 Figure 1906A-B: DNA328789, 344240.3, 214770.at  
 Figure 1907: PRO84534  
 Figure 1908A-B: DNA328790, 481415.9, 214786.at  
 Figure 1909: PRO84535  
 Figure 1910: DNA328791, 1383762.1, 214790.at  
 Figure 1911: PRO84536  
 Figure 1912: DNA328792, 7692351.10, 214830.at  
 Figure 1913: PRO84537  
 Figure 1914: DNA328314, BC022780, 214841.at  
 Figure 1915: PRO84182  
 Figure 1916: DNA83102, DNA83102, 214866.at  
 Figure 1917: PRO2591  
 Figure 1918: DNA161326, DNA161326, 214934.at  
 Figure 1919: DNA328794, 1099353.2, 214974.x.at  
 Figure 1920: PRO84539  
 Figure 1921: DNA328795, AF057354, 214975.s.at  
 Figure 1922: DNA328796, HSM800535, 215078.at  
 Figure 1923: DNA328797, 000092.6, 215087.at  
 Figure 1924: PRO84540  
 Figure 1925: DNA328798, NP\_002088.1, 215091.s.at  
 Figure 1926: PRO84541  
 Figure 1927: DNA328799, BC008376, 215101.s.at  
 Figure 1928: PRO1721  
 Figure 1929: DNA270522, NP\_006013.1, 215111.s.at  
 Figure 1930: PRO58899  
 Figure 1931: DNA328800, 194537.1, 215224.at  
 Figure 1932: PRO84542  
 Figure 1933A-B: DNA327827, HSM800826, 215235.at  
 Figure 1934A-B: DNA226905, NP\_055672.1, 215342.s.at  
 Figure 1935: PRO37368  
 Figure 1936: DNA327831, NP\_076956.1, 215380.s.at  
 Figure 1937: PRO83783  
 Figure 1938: DNA328801, 407831.1, 215392.at  
 Figure 1939: PRO84543  
 Figure 1940A-B: DNA328802, C6orf5, 215411.s.at  
 Figure 1941: PRO84544  
 Figure 1942: DNA275385, NP\_002085.1, 215438.x.at  
 Figure 1943: PRO63048  
 Figure 1944: DNA328803, BAA91443.1, 215440.s.at  
 Figure 1945: PRO84545  
 Figure 1946: DNA328804, 403621.1, 215767.at  
 Figure 1947: PRO84546  
 Figure 1948A-B: DNA328805, BAA86482.1, 215785.s.at  
 Figure 1949: PRO84547  
 Figure 1950: DNA328806, 208045.1, 216109.at  
 Figure 1951: PRO84548  
 Figure 1952: DNA269532, NP\_004802.1, 216250.s.at  
 Figure 1953: PRO57948  
 Figure 1954: DNA328807, AAH10129.1, 216483.s.at  
 Figure 1955: PRO84549  
 Figure 1956: DNA188349, NP\_002973.1, 216598.s.at  
 Figure 1957: PRO21884  
 Figure 1958: DNA328808, 1099517.2, 216607.s.at  
 Figure 1959: PRO84550  
 Figure 1960: DNA328809, PTPN12, 216915.s.at  
 Figure 1961: PRO4803  
 Figure 1962: DNA328810, NP\_001770.1, 216942.s.at  
 Figure 1963: PRO2557  
 Figure 1964A-C: DNA328811, NP\_002213.1, 216944.s.at  
 Figure 1965: PRO84551  
 Figure 1966: DNA328812, BAA86575.1, 216997.x.at  
 Figure 1967: PRO84552  
 Figure 1968A-B: DNA328813, BAA76774.1, 217118.s.at  
 Figure 1969: PRO84553  
 Figure 1970A-B: DNA328814, HUMMHHLAJC, 217436.x.at  
 Figure 1971A-B: DNA328815, 331104.2, 217521.at  
 Figure 1972: PRO84554  
 Figure 1973: DNA328816, 1446567.1, 217526.at  
 Figure 1974: PRO84555  
 Figure 1975A-B: DNA255619, AF054589, 217599.s.at  
 Figure 1976: PRO50682  
 Figure 1977: DNA327848, NP\_005998.1, 217649.at  
 Figure 1978: PRO83793  
 Figure 1979: DNA328817, 1498470.1, 217678.at  
 Figure 1980: PRO84556  
 Figure 1981: DNA328818, NP\_071435.1, 217730.at  
 Figure 1982: PRO38175  
 Figure 1983: DNA327935, NP\_079422.1, 217745.s.at  
 Figure 1984: PRO83866  
 Figure 1985A-B: DNA88040, NP\_000005.1, 217757.at  
 Figure 1986: PRO2632  
 Figure 1987A-B: DNA88226, NP\_000055.1, 217767.at  
 Figure 1988: PRO2237

- Figure 1989: DNA325821, NP\_057016.1, 217769.s.at  
 Figure 1990: PRO82287  
 Figure 1991: DNA227358, NP\_057479.1, 217777.s.at  
 Figure 1992: PRO37821  
 Figure 1993: DNA328819, NP\_057145.1, 217783.s.at  
 Figure 1994: PRO84557  
 Figure 1995: DNA327850, NP\_006546.1, 217785.s.at  
 Figure 1996: PRO60803  
 Figure 1997: DNA328830, NP\_056525.1, 217807.s.at  
 Figure 1998: PRO84173  
 Figure 1999: DNA328820, NP\_077022.1, 217808.s.at  
 Figure 2000: PRO84558  
 Figure 2001: DNA328821, NP\_006708.1, 217813.s.at  
 Figure 2002: PRO84559  
 Figure 2003: DNA328822, AK001511, 217830.s.at  
 Figure 2004: PRO84560  
 Figure 2005: DNA328823, NP\_057421.1, 217838.s.at  
 Figure 2006: PRO84561  
 Figure 2007: DNA226759, NP\_054775.1, 217845.x.at  
 Figure 2008: PRO37222  
 Figure 2009: DNA327939, NP\_060654.1, 217852.s.at  
 Figure 2010: PRO83869  
 Figure 2011A-B: DNA324921, NP\_073585.6, 217853.at  
 Figure 2012: PRO81523  
 Figure 2013: DNA328824, DREV1, 217868.s.at  
 Figure 2014: PRO84562  
 Figure 2015: DNA225604, NP\_057226.1, 217869.at  
 Figure 2016: PRO36067  
 Figure 2017: DNA326937, NP\_002406.1, 217871.s.at  
 Figure 2018: PRO83255  
 Figure 2019: DNA255145, NP\_060917.1, 217882.at  
 Figure 2020: PRO50225  
 Figure 2021A-B: DNA328825, 1398762.11, 217886.at  
 Figure 2022: PRO84563  
 Figure 2023: DNA189504, NP\_064539.1, 217898.at  
 Figure 2024: PRO25402  
 Figure 2025: DNA328826, NP\_004272.2, 217911.s.at  
 Figure 2026: PRO84564  
 Figure 2027: DNA328827, NP\_076869.1, 217949.s.at  
 Figure 2028: PRO21784  
 Figure 2029: DNA328828, NP\_067027.1, 217956.s.at  
 Figure 2030: PRO84565  
 Figure 2031: DNA328829, NP\_057230.1, 217959.s.at  
 Figure 2032: PRO84566  
 Figure 2033: DNA328830, NP\_061118.1, 217962.at  
 Figure 2034: PRO84567  
 Figure 2035: DNA327855, NP\_057387.1, 217975.at  
 Figure 2036: PRO83367  
 Figure 2037: DNA328831, NP\_057329.1, 217989.at  
 Figure 2038: PRO233  
 Figure 2039: DNA328832, NP\_067022.1, 217995.at  
 Figure 2040: PRO84568  
 Figure 2041: DNA328833, BC018929, 217996.at  
 Figure 2042: PRO84569  
 Figure 2043: DNA328834, AF220656, 217997.at  
 Figure 2044: DNA326005, NP\_057004.1, 218007.s.at  
 Figure 2045: PRO82446  
 Figure 2046: DNA328835, NP\_068760.1, 218019.s.at  
 Figure 2047: PRO84571  
 Figure 2048: DNA328836, NP\_054894.1, 218027.at  
 Figure 2049: PRO84572  
 Figure 2050: DNA328837, NP\_057149.1, 218046.s.at  
 Figure 2051: PRO81876  
 Figure 2052: DNA328838, NP\_054797.2, 218049.s.at  
 Figure 2053: PRO70319  
 Figure 2054: DNA328839, NP\_057180.1, 218059.at  
 Figure 2055: PRO84573  
 Figure 2056: DNA328840, NP\_060481.1, 218067.s.at  
 Figure 2057: PRO84574  
 Figure 2058: DNA328841, NP\_060557.2, 218073.s.at  
 Figure 2059: PRO84575  
 Figure 2060A-C: DNA328842, 235943.8, 218098.at  
 Figure 2061: PRO84576  
 Figure 2062: DNA328843, NP\_060939.1, 218099.at  
 Figure 2063: PRO84577  
 Figure 2064: DNA328844, NP\_061156.1, 218111.s.at  
 Figure 2065: PRO82111  
 Figure 2066: DNA227498, NP\_002125.3, 218120.s.at  
 Figure 2067: PRO37961  
 Figure 2068: DNA328845, NP\_060615.1, 218126.at  
 Figure 2069: PRO10274  
 Figure 2070: DNA227264, LOC51312, 218136.s.at  
 Figure 2071: PRO37727  
 Figure 2072: DNA327857, NP\_057386.1, 218142.s.at  
 Figure 2073: PRO83799  
 Figure 2074: DNA325852, NP\_078813.1, 218153.at  
 Figure 2075: PRO82314  
 Figure 2076: DNA328846, NP\_060522.2, 218169.at  
 Figure 2077: PRO84578  
 Figure 2078: DNA228094, NP\_079416.1, 218175.at  
 Figure 2079: PRO38557  
 Figure 2080: DNA328847, NP\_056338.1, 218194.at  
 Figure 2081: PRO84579  
 Figure 2082: DNA150593, NP\_054747.1, 218196.at  
 Figure 2083: PRO12353  
 Figure 2084: DNA256555, NP\_060042.1, 218205.s.at  
 Figure 2085: PRO51586  
 Figure 2086: DNA328848, NP\_004522.1, 218212.s.at  
 Figure 2087: PRO84580  
 Figure 2088: DNA271622, NP\_006020.3, 218224.at  
 Figure 2089: PRO59909  
 Figure 2090: DNA324353, NP\_004538.2, 218226.s.at  
 Figure 2091: PRO81026  
 Figure 2092: DNA328849, NP\_057075.1, 218232.at  
 Figure 2093: PRO4382  
 Figure 2094: DNA328850, NP\_057187.1, 218254.s.at  
 Figure 2095: PRO84581  
 Figure 2096: DNA273230, NP\_060914.1, 218273.s.at  
 Figure 2097: PRO61257  
 Figure 2098: DNA328851, NP\_068590.1, 218276.s.at  
 Figure 2099: PRO84582

Figure 2100: DNA323953, NP\_003507.1, 218280.x.at  
 Figure 2101: PRO80685  
 Figure 2102: DNA254824, AF267865, 218294.s.at  
 Figure 2103: PRO49920  
 Figure 2104A-B: DNA328852, NP\_003609.2, 218311.at  
 Figure 2105: PRO84583  
 Figure 2106A-B: DNA328853, NP\_065702.2, 218319.at  
 Figure 2107: PRO84584  
 Figure 2108: DNA328854, NP\_056979.1, 218350.s.at  
 Figure 2109: PRO84585  
 Figure 2110: DNA328855, NP\_076952.1, 218375.at  
 Figure 2111: PRO9771  
 Figure 2112: DNA328856, NP\_068376.1, 218380.at  
 Figure 2113: PRO84586  
 Figure 2114: DNA328857, NP\_037481.1, 218407.x.at  
 Figure 2115: PRO84587  
 Figure 2116: DNA324953, NP\_057412.1, 218412.s.at  
 Figure 2117: PRO81550  
 Figure 2118A-B: DNA255062, NP\_060704.1, 218424.s.at  
 Figure 2119: PRO50149  
 Figure 2120: DNA150661, NP\_057162.1, 218446.s.at  
 Figure 2121: PRO12398  
 Figure 2122: DNA326218, NP\_064573.1, 218447.at  
 Figure 2123: PRO82631  
 Figure 2124: DNA328858, HEBP1, 218450.at  
 Figure 2125: PRO84588  
 Figure 2126: DNA327942, NP\_060596.1, 218465.at  
 Figure 2127: PRO83870  
 Figure 2128: DNA328859, AF154054, 218468.s.at  
 Figure 2129: PRO1608  
 Figure 2130A-B: DNA328860, NP\_037504.1, 218469.at  
 Figure 2131: PRO1608  
 Figure 2132: DNA328861, NP\_057030.2, 218472.s.at  
 Figure 2133: PRO84589  
 Figure 2134: DNA328862, NP\_057626.2, 218499.at  
 Figure 2135: PRO84590  
 Figure 2136: DNA328863, NP\_060264.1, 218503.at  
 Figure 2137: PRO84591  
 Figure 2138: DNA328864, NP\_060726.2, 218512.at  
 Figure 2139: PRO84592  
 Figure 2140: DNA255432, NP\_060283.1, 218516.s.at  
 Figure 2141: PRO50499  
 Figure 2142: DNA194326, NP\_065713.1, 218538.s.at  
 Figure 2143: PRO23708  
 Figure 2144: DNA328865, NP\_064587.1, 218557.at  
 Figure 2145: PRO84593  
 Figure 2146: DNA328866, NP\_005691.1, 218567.x.at  
 Figure 2147: PRO69644  
 Figure 2148: DNA328867, NP\_085053.1, 218600.at  
 Figure 2149: PRO84594  
 Figure 2150: DNA328868, NP\_057629.1, 218611.at  
 Figure 2151: PRO84595  
 Figure 2152: DNA328869, NP\_060892.1, 218613.at  
 Figure 2153: PRO84596  
 Figure 2154: DNA328870, NP\_060639.1, 218614.at  
 Figure 2155: PRO84597  
 Figure 2156: DNA256870, NP\_073600.1, 218618.s.at  
 Figure 2157: PRO51800  
 Figure 2158: DNA254898, NP\_060840.1, 218627.at  
 Figure 2159: PRO49988  
 Figure 2160: DNA328871, NP\_068378.1, 218631.at  
 Figure 2161: PRO84598  
 Figure 2162: DNA328872, NP\_036528.1, 218634.at  
 Figure 2163: PRO84599  
 Figure 2164: DNA328873, NP\_057041.1, 218698.at  
 Figure 2165: PRO84600  
 Figure 2166: DNA324621, NP\_054754.1, 218705.s.at  
 Figure 2167: PRO1285  
 Figure 2168: DNA328874, NP\_054778.1, 218723.s.at  
 Figure 2169: PRO84601  
 Figure 2170: DNA328875, NP\_064554.2, 218729.at  
 Figure 2171: PRO84602  
 Figure 2172: DNA328876, NP\_060582.1, 218772.x.at  
 Figure 2173: PRO84603  
 Figure 2174: DNA328877, BC020507, 218821.at  
 Figure 2175: PRO84604  
 Figure 2176: DNA328878, NP\_060104.1, 218823.s.at  
 Figure 2177: PRO84605  
 Figure 2178: DNA328879, NP\_064570.1, 218845.at  
 Figure 2179: PRO84606  
 Figure 2180: DNA227367, NP\_062456.1, 218853.s.at  
 Figure 2181: PRO37830  
 Figure 2182: DNA327872, NP\_057713.1, 218856.at  
 Figure 2183: PRO83812  
 Figure 2184: DNA328880, NP\_060369.1, 218872.at  
 Figure 2185: PRO84607  
 Figure 2186: DNA328881, NP\_057706.1, 218890.x.at  
 Figure 2187: PRO49469  
 Figure 2188: DNA287166, NP\_055129.1, 218943.s.at  
 Figure 2189: PRO69459  
 Figure 2190: DNA328882, NP\_109589.1, 218967.s.at  
 Figure 2191: PRO61822  
 Figure 2192: DNA327211, NP\_075053.1, 218989.x.at  
 Figure 2193: PRO71052  
 Figure 2194: DNA255929, NP\_060935.1, 218992.at  
 Figure 2195: PRO50982  
 Figure 2196: DNA328883, NP\_037474.1, 218996.at  
 Figure 2197: PRO84608  
 Figure 2198: DNA227194, FLJ11000, 218999.at  
 Figure 2199: PRO37657  
 Figure 2200: DNA328884, NP\_054884.1, 219006.at  
 Figure 2201: PRO84609  
 Figure 2202: DNA227187, NP\_057703.1, 219014.at  
 Figure 2203: PRO37650  
 Figure 2204: DNA328885, NP\_061108.2, 219017.at  
 Figure 2205: PRO50294  
 Figure 2206A-B: DNA255239, NP\_004832.1, 219026.s.at

- Figure 2207: PRO50316  
 Figure 2208: DNA328886, NP\_078811.1, 219040.at  
 Figure 2209: PRO84610  
 Figure 2210: DNA328887, NP\_061907.1, 219045.at  
 Figure 2211: PRO84611  
 Figure 2212: DNA328888, NP\_060436.1, 219053.s.at  
 Figure 2213: PRO84612  
 Figure 2214: DNA328889, NP\_006005.1, 219061.s.at  
 Figure 2215: PRO84613  
 Figure 2216: DNA328890, NP\_060403.1, 219093.at  
 Figure 2217: PRO84614  
 Figure 2218: DNA327877, NP\_065108.1, 219099.at  
 Figure 2219: PRO83816  
 Figure 2220: DNA328891, NP\_060263.1, 219143.s.at  
 Figure 2221: PRO84615  
 Figure 2222: DNA210216, NP\_006860.1, 219150.s.at  
 Figure 2223: PRO33752  
 Figure 2224: DNA328892, NP\_067643.2, 219165.at  
 Figure 2225: PRO84616  
 Figure 2226A-B: DNA328893, NP\_065699.1, 219201.s.at  
 Figure 2227: PRO9914  
 Figure 2228: DNA287235, NP\_060598.1, 219204.s.at  
 Figure 2229: PRO69514  
 Figure 2230: DNA225594, NP\_037404.1, 219229.at  
 Figure 2231: PRO36057  
 Figure 2232: DNA328894, NP\_060796.1, 219243.at  
 Figure 2233: PRO84617  
 Figure 2234: DNA328895, NP\_071762.2, 219259.at  
 Figure 2235: PRO1317  
 Figure 2236: DNA328896, NP\_079037.1, 219265.at  
 Figure 2237: PRO84618  
 Figure 2238: DNA328897, TRPV2, 219282.s.at  
 Figure 2239: PRO12382  
 Figure 2240: DNA273489, NP\_055210.1, 219290.x.at  
 Figure 2241: PRO61472  
 Figure 2242A-B: DNA328898, NP\_060261.1, 219316.s.at  
 Figure 2243: PRO84619  
 Figure 2244: DNA328899, NP\_061024.1, 219326.s.at  
 Figure 2245: PRO84620  
 Figure 2246A-B: DNA255889, NP\_061764.1, 219340.s.at  
 Figure 2247: PRO50942  
 Figure 2248: DNA328900, NP\_060814.1, 219344.at  
 Figure 2249: PRO84621  
 Figure 2250: DNA254518, NP\_057354.1, 219371.s.at  
 Figure 2251: PRO49625  
 Figure 2252: DNA188342, NP\_064510.1, 219385.at  
 Figure 2253: PRO21718  
 Figure 2254: DNA256417, NP\_077271.1, 219402.s.at  
 Figure 2255: PRO51457  
 Figure 2256A-B: DNA327887, NP\_006656.1, 219403.s.at  
 Figure 2257: PRO83823  
 Figure 2258: DNA327888, NP\_071732.1, 219412.at  
 Figure 2259: PRO83824  
 Figure 2260: DNA328901, FLJ20533, 219449.s.at  
 Figure 2261: PRO84622  
 Figure 2262: DNA328902, NP\_071750.1, 219452.at  
 Figure 2263: PRO84623  
 Figure 2264: DNA328903, NP\_002805.1, 219485.s.at  
 Figure 2265: PRO84624  
 Figure 2266: DNA328904, NP\_076941.1, 219491.at  
 Figure 2267: PRO84625  
 Figure 2268A-B: DNA328905, NP\_075392.1, 219496.at  
 Figure 2269: PRO84626  
 Figure 2270: DNA328906, NP\_078855.1, 219506.at  
 Figure 2271: PRO84627  
 Figure 2272: DNA328907, NP\_000067.1, 219534.x.at  
 Figure 2273: PRO84628  
 Figure 2274: DNA328908, 7691567.2, 219540.at  
 Figure 2275: PRO84629  
 Figure 2276: DNA225636, NP\_065696.1, 219557.s.at  
 Figure 2277: PRO36099  
 Figure 2278A-B: DNA328909, NP\_078800.2, 219558.at  
 Figure 2279: PRO84630  
 Figure 2280: DNA328910, NP\_057666.1, 219593.at  
 Figure 2281: PRO38848  
 Figure 2282: DNA328911, MS4A4A, 219607.s.at  
 Figure 2283: PRO84631  
 Figure 2284: DNA328912, NP\_060287.1, 219622.at  
 Figure 2285: PRO84632  
 Figure 2286: DNA328913, NP\_079213.1, 219631.at  
 Figure 2287: PRO84633  
 Figure 2288: DNA328914, NP\_060883.1, 219634.at  
 Figure 2289: PRO36664  
 Figure 2290: DNA327892, NP\_060470.1, 219648.at  
 Figure 2291: PRO83828  
 Figure 2292: DNA328915, NP\_055056.2, 219654.at  
 Figure 2293: PRO84634  
 Figure 2294: DNA228002, NP\_071744.1, 219666.at  
 Figure 2295: PRO38465  
 Figure 2296: DNA328916, NP\_071932.1, 219678.x.at  
 Figure 2297: PRO84635  
 Figure 2298: DNA287206, NP\_060124.1, 219691.at  
 Figure 2299: PRO69488  
 Figure 2300: DNA328917, NP\_061838.1, 219725.at  
 Figure 2301: PRO7306  
 Figure 2302: DNA328918, NP\_078935.1, 219770.at  
 Figure 2303: PRO84636  
 Figure 2304: DNA328919, NP\_078987.1, 219777.at  
 Figure 2305: PRO84637  
 Figure 2306: DNA227152, NP\_038467.1, 219788.at  
 Figure 2307: PRO37615  
 Figure 2308: DNA328920, NP\_061129.1, 219837.s.at  
 Figure 2309: PRO4425  
 Figure 2310: DNA256033, NP\_060164.1, 219858.s.at  
 Figure 2311: PRO51081  
 Figure 2312: DNA254838, NP\_078904.1, 219874.at

- Figure 2313: PRO49933  
Figure 2314: DNA328921, NP\_057079.1, 219878.s\_at  
Figure 2315: PRO84638  
Figure 2316: DNA256325, NP\_005470.1, 219889.at  
Figure 2317: PRO51367  
Figure 2318: DNA328922, NP\_037384.1, 219890.at  
Figure 2319: PRO84639  
Figure 2320: DNA328923, NP\_075379.1, 219892.at  
Figure 2321: PRO84640  
Figure 2322: DNA256608, NP\_060408.1, 219895.at  
Figure 2323: PRO51611  
Figure 2324: DNA328924, NP\_057150.2, 219933.at  
Figure 2325: PRO84641  
Figure 2326: DNA255456, NP\_057268.1, 219947.at  
Figure 2327: PRO50523  
Figure 2328: DNA227804, NP\_065394.1, 219952.s\_at  
Figure 2329: PRO38267  
Figure 2330: DNA328925, NP\_076403.1, 220005.at  
Figure 2331: PRO84642  
Figure 2332: DNA256467, NP\_079054.1, 220009.at  
Figure 2333: PRO51504  
Figure 2334A-B: DNA292946, NP\_061160.1, 220023.at  
Figure 2335: PRO70613  
Figure 2336: DNA171414, NP\_009130.1, 220034.at  
Figure 2337: PRO20142  
Figure 2338: DNA328926, NP\_064703.1, 220046.s\_at  
Figure 2339: PRO84643  
Figure 2340A-B: DNA221079, NP\_071445.1, 220066.at  
Figure 2341: PRO34753  
Figure 2342: DNA256091, NP\_071385.1, 220094.s\_at  
Figure 2343: PRO51141  
Figure 2344: DNA328927, NP\_078993.2, 220122.at  
Figure 2345: PRO84644  
Figure 2346: DNA328928, NP\_068377.1, 220178.at  
Figure 2347: PRO84645  
Figure 2348: DNA324716, NP\_463459.1, 220189.s\_at  
Figure 2349: PRO81347  
Figure 2350: DNA228059, NP\_073742.1, 220199.s\_at  
Figure 2351: PRO38522  
Figure 2352: DNA328929, NP\_060375.1, 220240.s\_at  
Figure 2353: PRO84646  
Figure 2354A-B: DNA328930, NP\_038465.1, 220253.s\_at  
Figure 2355: PRO23525  
Figure 2356: DNA328931, NP\_004226.1, 220266.s\_at  
Figure 2357: PRO84647  
Figure 2358: DNA328932, NP\_079057.1, 220301.at  
Figure 2359: PRO84648  
Figure 2360: DNA328933, NP\_057466.1, 220307.at  
Figure 2361: PRO9891  
Figure 2362: DNA256735, NP\_060175.1, 220333.at  
Figure 2363: PRO51669  
Figure 2364A-B: DNA328934, EML4, 220386.s\_at  
Figure 2365: PRO84649  
Figure 2366: DNA328935, NP\_009002.1, 220387.s\_at  
Figure 2367: PRO84650  
Figure 2368: DNA254861, MCOLN3, 220484.at  
Figure 2369: PRO49953  
Figure 2370: DNA328936, NP\_066998.1, 220491.at  
Figure 2371: PRO1003  
Figure 2372: DNA328937, PHEMX, 220558.x\_at  
Figure 2373: PRO12380  
Figure 2374: DNA328938, NP\_060617.1, 220643.s\_at  
Figure 2375: PRO84651  
Figure 2376: DNA323756, NP\_057267.2, 220688.s\_at  
Figure 2377: PRO80512  
Figure 2378: DNA328939, NP\_008834.1, 220741.s\_at  
Figure 2379: PRO84652  
Figure 2380: DNA288247, NP\_478059.1, 220892.s\_at  
Figure 2381: PRO70011  
Figure 2382: DNA328940, NP\_078893.1, 220933.s\_at  
Figure 2383: PRO84653  
Figure 2384: DNA328941, NP\_055218.2, 220937.s\_at  
Figure 2385: PRO84654  
Figure 2386: DNA327953, NP\_055182.2, 220942.x\_at  
Figure 2387: PRO83878  
Figure 2388A-B: DNA323882, NP\_000692.2, 220948.s\_at  
Figure 2389: PRO80625  
Figure 2390: DNA327917, NP\_112240.1, 220966.x\_at  
Figure 2391: PRO83852  
Figure 2392: DNA328942, NP\_112216.2, 220985.s\_at  
Figure 2393: PRO84655  
Figure 2394: DNA328943, NP\_036566.1, 221041.s\_at  
Figure 2395: PRO51680  
Figure 2396: DNA328944, NP\_060554.1, 221078.s\_at  
Figure 2397: PRO84656  
Figure 2398: DNA328945, NP\_079177.2, 221081.s\_at  
Figure 2399: PRO84657  
Figure 2400: DNA328946, NP\_055164.1, 221087.s\_at  
Figure 2401: PRO12343  
Figure 2402: DNA328947, NP\_055245.1, 221188.s\_at  
Figure 2403: PRO84658  
Figure 2404: DNA257293, NP\_110396.1, 221210.s\_at  
Figure 2405: PRO51888  
Figure 2406: DNA327920, NP\_110431.1, 221245.s\_at  
Figure 2407: PRO83855  
Figure 2408A-C: DNA328287, NP\_072174.2, 221246.x\_at  
Figure 2409: PRO84163  
Figure 2410: DNA328948, NP\_110437.1, 221253.s\_at  
Figure 2411: PRO84659  
Figure 2412: DNA256432, NP\_110415.1, 221266.s\_at  
Figure 2413: PRO51471  
Figure 2414: DNA328027, NP\_112570.2, 221437.s\_at  
Figure 2415: PRO83944  
Figure 2416A-B: DNA272014, AF084555, 221482.s\_at  
Figure 2417: PRO60289  
Figure 2418: DNA328949, AF157510, 221487.s\_at

- Figure 2419: PRO84660  
Figure 2420: DNA328950, NP\_057025.1, 221504.s.at  
Figure 2421: PRO84661  
Figure 2422A-B: DNA328951, HSM802232, 221523.s.at  
Figure 2423: PRO84662  
Figure 2424: DNA328952, NP\_067067.1, 221524.s.at  
Figure 2425: PRO84663  
Figure 2426A-B: DNA273901, NP\_110389.1, 221530.s.at  
Figure 2427: PRO61855  
Figure 2428: DNA274676, DKFZp564A176Homo, 221538.s.at  
Figure 2429: DNA328953, NP\_004086.1, 221539.at  
Figure 2430: PRO70296  
Figure 2431A-B: DNA328954, NP\_113664.1, 221541.at  
Figure 2432: PRO9851  
Figure 2433A-B: DNA269992, HUMACYLCOA, 221561.at  
Figure 2434: PRO58388  
Figure 2435: DNA328955, NP\_054887.1, 221570.s.at  
Figure 2436: PRO84664  
Figure 2437A-B: DNA328956, AF110908, 221571.at  
Figure 2438: DNA188321, NP\_004855.1, 221577.x.at  
Figure 2439: PRO21896  
Figure 2440: DNA328957, WBSCR5, 221581.s.at  
Figure 2441: PRO23859  
Figure 2442: DNA328958, BC001663, 221593.s.at  
Figure 2443: PRO84665  
Figure 2444: DNA328959, NP\_077027.1, 221620.s.at  
Figure 2445: PRO4302  
Figure 2446: DNA254777, NP\_055140.1, 221676.s.at  
Figure 2447: PRO49875  
Figure 2448: DNA327526, NP\_065727.2, 221679.s.at  
Figure 2449: PRO83574  
Figure 2450: DNA328960, NP\_076426.1, 221692.s.at  
Figure 2451: PRO84666  
Figure 2452: DNA327929, AK001785, 221748.s.at  
Figure 2453: PRO83861  
Figure 2454: DNA328961, NP\_443112.1, 221756.at  
Figure 2455: PRO84667  
Figure 2456: DNA328962, BC021574, 221759.at  
Figure 2457: PRO82746  
Figure 2458A-B: DNA328963, 328765.9, 221760.at  
Figure 2459: PRO84668  
Figure 2460A-B: DNA327930, 1455324.9, 221765.at  
Figure 2461: PRO83862  
Figure 2462: DNA328964, AK056028, 221770.at  
Figure 2463: PRO84669  
Figure 2464A-C: DNA328965, AB051505, 221778.at  
Figure 2465A-B: DNA328966, BAB14908.1, 221790.s.at  
Figure 2466: PRO84670  
Figure 2467: DNA328967, BC017905, 221815.at  
Figure 2468: PRO84671  
Figure 2469: DNA274058, NP\_057203.1, 221816.s.at  
Figure 2470: PRO61999  
Figure 2471A-B: DNA328968, 1322249.6, 221830.at  
Figure 2472: PRO62511  
Figure 2473: DNA272419, AF105036, 221841.s.at  
Figure 2474: PRO60672  
Figure 2475: DNA299882, DNA299882, 221872.at  
Figure 2476: PRO70856  
Figure 2477: DNA328969, 334394.2, 221878.at  
Figure 2478: PRO84672  
Figure 2479: DNA327933, 1452741.11, 221899.at  
Figure 2480: PRO83865  
Figure 2481: DNA328970, NP\_057696.1, 221920.s.at  
Figure 2482: PRO84673  
Figure 2483: DNA328971, AK000472, 221923.s.at  
Figure 2484: PRO84674  
Figure 2485: DNA254787, AK023140, 221935.s.at  
Figure 2486: PRO49885  
Figure 2487: DNA327114, NP\_006004.1, 221989.at  
Figure 2488: PRO62466  
Figure 2489: DNA328972, BC009950, 222001.x.at  
Figure 2490: DNA328973, NP\_115538.1, 222024.s.at  
Figure 2491: PRO82497  
Figure 2492: DNA119482, DNA119482, 222108.at  
Figure 2493: PRO9850  
Figure 2494: DNA328974, NP\_061893.1, 222116.s.at  
Figure 2495: PRO84676  
Figure 2496: DNA287209, NP\_056350.1, 222154.s.at  
Figure 2497: PRO69490  
Figure 2498: DNA328975, NP\_078807.1, 222155.s.at  
Figure 2499: PRO47688  
Figure 2500: DNA328976, BC019091, 222206.s.at  
Figure 2501: PRO84677  
Figure 2502: DNA256784, NP\_075069.1, 222209.s.at  
Figure 2503: PRO51716  
Figure 2504: DNA328977, NP\_071344.1, 222216.s.at  
Figure 2505: PRO84678  
Figure 2506: DNA328978, NP\_060373.1, 222244.s.at  
Figure 2507: PRO84679  
Figure 2508A-B: DNA328979, 006242.19, 222266.at  
Figure 2509: PRO84680  
Figure 2510: DNA328980, 7692031.1, 222273.at  
Figure 2511: PRO84681  
Figure 2512: DNA328981, AF443871, 222294.s.at  
Figure 2513: PRO24633  
Figure 2514: DNA328982, 154391.1, 222313.at  
Figure 2515: PRO84682  
Figure 2516: DNA328983, 206335.1, 222366.at  
Figure 2517: PRO84683

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

### I. Definitions

The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the "signal-peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

"PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length,



alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

"Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X", "Y" and "Z" each represent different hypothetical amino acid residues.

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-

BLAST-2 computer program (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an the amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

"PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid

sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alternatively at least about 240 nucleotides in length, alternatively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, California or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence

D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction  $W/Z$

5

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span = 1, overlap fraction = 0.125, word threshold (T) = 11, and scoring matrix = BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, MD. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask = yes, strand = all, expected occurrences = 10, minimum low complexity length = 15/5, multi-pass e-value = 0.01, constant for multi-pass = 25, dropoff for final gapped alignment = 25 and scoring matrix = BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

5

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-PRO monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polypeptidic specificity, single chain anti-PRO antibodies, and fragments of anti-PRO antibodies (see below). The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a

solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate); 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

"Active" or "activity" for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those

in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time.

5 "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

10 Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of  
15 physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating  
20 agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata et al., *Protein Eng.* 8(10): 1057-1062 [1995]); single-chain  
25 antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

30 "Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V<sub>H</sub>-V<sub>L</sub> dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising  
35 only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge  
40 region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains



bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

"Single-chain Fv" or "sFv" antibody fragments comprise the V<sub>H</sub> and V<sub>L</sub> domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) in the same polypeptide chain (V<sub>H</sub>-V<sub>L</sub>). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass

(e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Patent No.

5 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

10 A "small molecule" is defined herein to have a molecular weight below about 500 Daltons.

The term "immune related disease" means a disease in which a component of the immune system of a mammal causes, mediates or otherwise contributes to a morbidity in the mammal. Also included are diseases in which stimulation or intervention of the immune response has an ameliorative effect on progression of the disease. Included within this term are immune-mediated inflammatory diseases, non-immune-mediated inflammatory diseases, infectious diseases, immunodeficiency diseases, neoplasia, etc.

15 The term "monocyte/macrophage mediated disease" means a disease in which monocytes/macrophages directly or indirectly mediate or otherwise contribute to a morbidity in a mammal. The monocyte/macrophage mediated disease may be associated with cell mediated effects, lymphokine mediated effects, etc., and even effects associated with other immune cells if the cells are stimulated, for example, by the lymphokines secreted by monocytes/macrophages.

Examples of immune-related and inflammatory diseases, some of which are immune mediated, which can be treated according to the invention include systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, spondyloarthropathies, systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjögren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia (immune pancytopenia, paroxysmal nocturnal hemoglobinuria), autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis), demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory bowel disease (ulcerative colitis: Crohn's disease), gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft -versus-host-disease. Infectious diseases including viral diseases such as AIDS (HIV infection), hepatitis A, B, C, D, and E, herpes, etc., bacterial infections, fungal infections, protozoal infections and parasitic infections.

The term "effective amount" is a concentration or amount of a PRO polypeptide and/or agonist/antagonist which results in achieving a particular stated purpose. An "effective amount" of a PRO polypeptide or agonist or antagonist thereof may be determined empirically. Furthermore, a "therapeutically effective amount" is a concentration or amount of a PRO polypeptide and/or agonist/antagonist which is effective for achieving a stated therapeutic effect. This amount may also be determined empirically.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g.,  $I^{131}$ ,  $I^{125}$ ,  $Y^{90}$  and  $Re^{186}$ ), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include adriamycin, doxorubicin, epirubicin, 5-fluorouracil, cytosine arabinoside ("Ara-C"), cyclophosphamide, thiotepa, busulfan, cytoxin, taxoids, e.g., paclitaxel (Taxol, Bristol-Myers Squibb Oncology, Princeton, NJ), and doxetaxel (Taxotere, Rhône-Poulenc Rorer, Antony, France), toxotere, methotrexate, cisplatin, melphalan, vinblastine, bleomycin, etoposide, ifosfamide, mitomycin C, mitoxantrone, vincristine, vinorelbine, carboplatin, teniposide, daunomycin, carminomycin, aminopterin, dactinomycin, mitomycins, esperamicins (see U.S. Pat. No. 4,675,187), melphalan and other related nitrogen mustards. Also included in this definition are hormonal agents that act to regulate or inhibit hormone action on tumors such as tamoxifen and onapristone.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially cancer cell overexpressing any of the genes identified herein, either *in vitro* or *in vivo*. Thus, the growth inhibitory agent is one which significantly reduces the percentage of cells overexpressing such genes in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogens, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13.

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- $\alpha$  and - $\beta$ ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF- $\beta$ ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- $\alpha$  and TGF- $\beta$ ; insulin-like growth factor-I and -II; erythropoietin (EPO);

osteoinductive factors; interferons such as interferon- $\alpha$ , - $\beta$ , and - $\gamma$ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 $\alpha$ , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; a tumor necrosis factor such as TNF- $\alpha$  or TNF- $\beta$ ; and other polypeptide factors including LIF and kit ligand (KL).

5 As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the  
10 desired binding specificity which is other than the antigen recognition and binding site of an antibody (*i.e.*, is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE,  
15 IgD or IgM.

**Table 1**

```

/*
5  *
  * C-C increased from 12 to 15
  * Z is average of EQ
  * B is average of ND
  * match with stop is _M; stop-stop = 0; J (joker) match = 0
10 */
#define _M      -8      /* value of a match with a stop */

int  _day[26][26] = {
/*  A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
15 /* A */ { 2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ { 0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ { 0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ { 0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
20 /* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ { 1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
25 /* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ { 0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ { _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
30 0, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ { 1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ { 0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ { 1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
35 /* T */ { 1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ { 0, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ { 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
40 /* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ { 0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};

```

45

50

55

Table 1 (cont')

```

/*
*/
#include <stdio.h>
5  #include <ctype.h>

#define MAXJMP      16      /* max jumps in a diag */
#define MAXGAP      24      /* don't continue to penalize gaps larger than this */
#define JMPS        1024    /* max jmps in an path */
10  #define MX        4      /* save if there's at least MX-1 bases since last jmp */

#define DMAT        3      /* value of matching bases */
#define DMIS        0      /* penalty for mismatched bases */
#define DINS0       8      /* penalty for a gap */
15  #define DINS1     1      /* penalty per base */
#define PINS0       8      /* penalty for a gap */
#define PINS1       4      /* penalty per residue */

struct jmp {
20      short      n[MAXJMP]; /* size of jmp (neg for dely) */
      unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
}; /* limits seq to 2^16 -1 */

struct diag {
25      int      score; /* score at last jmp */
      long      offset; /* offset of prev block */
      short     ijmp; /* current jmp index */
      struct jmp jp; /* list of jmps */
30 };

struct path {
      int      spc; /* number of leading spaces */
      short     n[JMPS]; /* size of jmp (gap) */
35      int      x[JMPS]; /* loc of jmp (last elem before gap) */
};

char      *ofile; /* output file name */
char      *namex[2]; /* seq names: getseqs() */
40  char      *prog; /* prog name for err msgs */
char      *seqx[2]; /* seqs: getseqs() */
int      dmax; /* best diag: nw() */
int      dmax0; /* final diag */
int      dna; /* set if dna: main() */
int      endgaps; /* set if penalizing end gaps */
45  int      gapx, gapy; /* total gaps in seqs */
int      len0, len1; /* seq lens */
int      ngapx, ngapy; /* total size of gaps */
int      smax; /* max score: nw() */
int      *xbm; /* bitmap for matching */
50  long     offset; /* current offset in jmp file */
struct     diag *dx; /* holds diagonals */
struct     path pp[2]; /* holds path for seqs */

char      *calloc(), *malloc(), *index(), *strcpy();
55  char      *getseq(), *g_malloc();

```

Table 1 (cont')

```

/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
5  * where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case and may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
10  * Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
15 #include "nw.h"
#include "day.h"

static _dbval[26] = {
20 1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};

static _pbval[26] = {
25 1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4, 8, 16, 32, 64,
128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};

main(ac, av)
30 main
int ac;
char *av[ ];
{
35 prog = av[0];
if (ac != 3) {
fprintf(stderr, "usage: %s file1 file2\n", prog);
fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
fprintf(stderr, "The sequences can be in upper- or lower-case\n");
40 fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
fprintf(stderr, "Output is in the file \"align.out\"\n");
exit(1);
}
namex[0] = av[1];
namex[1] = av[2];
45 seqx[0] = getseq(namex[0], &len0);
seqx[1] = getseq(namex[1], &len1);
xbm = (dna)? _dbval : _pbval;

endgaps = 0; /* 1 to penalize endgaps */
50 ofile = "align.out"; /* output file */

nw(); /* fill in the matrix, get the possible jumps */
readjumps(); /* get the actual jumps */
print(); /* print stats, alignment */
55 cleanup(); /* unlink any tmp files */
}

```

**Table 1 (cont')**

```

/* do the alignment, return best score: main()
 * dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
 * pro: PAM 250 values
5  * When scores are equal, we prefer mismatches to any gap, prefer
   * a new gap to extending an ongoing gap, and prefer a gap in seqx
   * to a gap in seq y.
   */
nw()
10 {
    char      *px, *py;      /* seqs and ptrs */
    int        *ndely, *dely; /* keep track of dely */
    int        ndelx, delx;   /* keep track of delx */
15    int        *tmp;        /* for swapping row0, row1 */
    int        mis;          /* score for each type */
    int        ins0, ins1;    /* insertion penalties */
    register   id;           /* diagonal index */
    register   ij;           /* jump index */
20    register   *col0, *col1; /* score for curr, last row */
    register   xx, yy;       /* index into seqs */

    dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));

25    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
30    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;

    smax = -10000;
    if (endgaps) {
35        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;      /* Waterman Bull Math Biol 84 */
40    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;

    /* fill in match matrix
    */
45    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
50            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0] - ins1;
            ndelx = xx;
55        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
60        }
    }

```



**Table 1 (cont')**

...nw

```

5      for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
        mis = col0[yy-1];
        if (dna)
            mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
        else
            mis += _day[*px-'A'][*py-'A'];
10      /* update penalty for del in x seq;
        * favor new del over ongong del
        * ignore MAXGAP if weighting endgaps
        */
15      if (endgaps || ndely[yy] < MAXGAP) {
        if (col0[yy] - ins0 >= dely[yy]) {
            dely[yy] = col0[yy] - (ins0+ins1);
            ndely[yy] = 1;
        } else {
            dely[yy] -= ins1;
            ndely[yy]++;
20      }
        } else {
            if (col0[yy] - (ins0+ins1) >= dely[yy]) {
25              dely[yy] = col0[yy] - (ins0+ins1);
              ndely[yy] = 1;
            } else
                ndely[yy]++;
        }
30      /* update penalty for del in y seq;
        * favor new del over ongong del
        */
        if (endgaps || ndelx < MAXGAP) {
35          if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
          } else {
            delx -= ins1;
            ndelx++;
40          }
        } else {
            if (col1[yy-1] - (ins0+ins1) >= delx) {
45              delx = col1[yy-1] - (ins0+ins1);
              ndelx = 1;
            } else
                ndelx++;
        }
50      /* pick the maximum score; we're favoring
        * mis over any del and delx over dely
        */
55
60

```

Table 1 (cont')

...nw

```

5      id = xx - yy + len1 - 1;
      if (mis >= delx && mis >= dely[yy])
          coll[yy] = mis;
      else if (delx >= dely[yy]) {
          coll[yy] = delx;
          ij = dx[id].ijmp;
10      if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
          && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
          dx[id].ijmp++;
          if (++ij >= MAXJMP) {
15              writejumps(id);
              ij = dx[id].ijmp = 0;
              dx[id].offset = offset;
              offset += sizeof(struct jmp) + sizeof(offset);
          }
          dx[id].jp.n[ij] = ndelx;
          dx[id].jp.x[ij] = xx;
          dx[id].score = delx;
      }
      else {
25          coll[yy] = dely[yy];
          ij = dx[id].ijmp;
          if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
          && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
          dx[id].ijmp++;
          if (++ij >= MAXJMP) {
30              writejumps(id);
              ij = dx[id].ijmp = 0;
              dx[id].offset = offset;
              offset += sizeof(struct jmp) + sizeof(offset);
          }
          dx[id].jp.n[ij] = -ndely[yy];
          dx[id].jp.x[ij] = xx;
          dx[id].score = dely[yy];
      }
40      if (xx == len0 && yy < len1) {
          /* last col
          */
          if (endgaps)
              coll[yy] -= ins0+ins1*(len1-yy);
          if (coll[yy] > smax) {
45              smax = coll[yy];
              dmax = id;
          }
      }
50      if (endgaps && xx < len0)
          coll[yy-1] -= ins0+ins1*(len0-xx);
          if (coll[yy-1] > smax) {
              smax = coll[yy-1];
              dmax = id;
55      }
      tmp = col0; col0 = coll; coll = tmp;
  }
  (void) free((char *)ndely);
  (void) free((char *)dely);
  (void) free((char *)col0);
  (void) free((char *)col1);
60      }

```

Table 1 (cont')

```

/*
 *
 * print() -- only routine visible outside this module
5  *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[ ]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
10 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */
15
#include "nw.h"

#define SPC      3
#define P_LINE   256 /* maximum output line */
20 #define P_SPC   3    /* space between name or num and seq */

extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */
25

print()
{
    print
    {
        int lx, ly, firstgap, lastgap; /* overlap */
30
        if ((fx = fopen(ofile, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, ofile);
            cleanup(1);
        }
35
        fprintf(fx, "< first sequence: %s (length = %d)\n", namex[0], len0);
        fprintf(fx, "< second sequence: %s (length = %d)\n", namex[1], len1);
        olen = 60;
        lx = len0;
        ly = len1;
40
        firstgap = lastgap = 0;
        if (dmax < len1 - 1) { /* leading gap in x */
            pp[0].spc = firstgap = len1 - dmax - 1;
            ly -= pp[0].spc;
        }
45
        else if (dmax > len1 - 1) { /* leading gap in y */
            pp[1].spc = firstgap = dmax - (len1 - 1);
            lx -= pp[1].spc;
        }
        if (dmax0 < len0 - 1) { /* trailing gap in x */
50
            lastgap = len0 - dmax0 - 1;
            lx -= lastgap;
        }
        else if (dmax0 > len0 - 1) { /* trailing gap in y */
55
            lastgap = dmax0 - (len0 - 1);
            ly -= lastgap;
        }
        getmat(lx, ly, firstgap, lastgap);
        pr_align();
    }
60
}

```

**Table 1 (cont')**

```

/*
 * trace back the best path, count matches
 */
5 static
  getmat(lx, ly, firstgap, lastgap)
      int      lx, ly;
      int      firstgap, lastgap;
      /* "core" (minus endgaps) */
      /* leading trailing overlap */
10 {
    int      nm, i0, i1, siz0, siz1;
    char      outx[32];
    double    pct;
    register  n0, n1;
    register char *p0, *p1;
15
    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;
20
    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
30
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
35
        }
        else {
            if (xbm[*p0-'A']&xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
40
        }
45
    }

    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
55
        lx = (lx < ly)? lx : ly;
    pct = 100.*(double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "< %d match%s in an overlap of %d: %.2f percent similarity\n",
60
        nm, (nm == 1)? "" : "es", lx, pct);

```

getmat

**Table 1 (cont')**

```

fprintf(fx, "<gaps in first sequence: %d", gapx);
...getmat
5  if (gapx) {
    (void) sprintf(outx, "(%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);

10  fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);

15  }
    if (dna)
        fprintf(fx,
            "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
20  else
        fprintf(fx,
            "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
25  fprintf(fx,
            "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
    else
30  fprintf(fx, "<endgaps not penalized\n");
}
static      nm;          /* matches in core -- for checking */
static      lmax;        /* lengths of stripped file names */
static      ij[2];       /* jmp index for a path */
35  static    nc[2];       /* number at start of current line */
static      ni[2];       /* current elem number -- for gapping */
static      siz[2];
static char  *ps[2];      /* ptr to current element */
static char  *po[2];      /* ptr to next output char slot */
40  static char out[2][P_LINE]; /* output line */
static char  star[P_LINE]; /* set by stars() */

/*
 * print alignment of described in struct path pp[ ]
 */
45  static
pr_align()
    pr_align
{
50  int      nn;          /* char count */
    int      more;
    register i;

    for (i = 0, lmax = 0; i < 2; i++) {
55  nn = stripname(namex[i]);
        if (nn > lmax)
            lmax = nn;

        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
60  }
}

```

**Table 1 (cont')**

```

for (nn = nm = 0, more = 1; more;) {
...pr_align
    for (i = more = 0; i < 2; i++) {
5         /*
           * do we have more of this sequence?
           */
           if (!*ps[i])
10                continue;

           more++;

           if (pp[i].spc) { /* leading space */
15                 *po[i]++ = ' ';
                 pp[i].spc--;
           }
           else if (siz[i]) { /* in a gap */
20                 *po[i]++ = '-';
                 siz[i]--;
           }
           else { /* we're putting a seq element
                   */
25                 *po[i] = *ps[i];
                 if (islower(*ps[i]))
                     *ps[i] = toupper(*ps[i]);
                 po[i]++;
                 ps[i]++;

30                 /*
                   * are we at next gap for this seq?
                   */
                 if (ni[i] == pp[i].x[ij[i]]) {
40                     /*
                       * we need to merge all gaps
                       * at this location
                       */
                       siz[i] = pp[i].n[ij[i]++];
                       while (ni[i] == pp[i].x[ij[i]])
                           siz[i] += pp[i].n[ij[i]++];
                       }
                       ni[i]++;
                   }
           }
45         if (++nn == olen || !more && nn) {
                dumpblock();
                for (i = 0; i < 2; i++)
                    po[i] = out[i];
                nn = 0;
           }
50     }
}

/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
55 static
dumpblock()
{
    dumpblock
{
60     register i;
        for (i = 0; i < 2; i++)
            *po[i] = '\0';
    }
}

```

Table 1 (cont')

```

5      (void) putc('\n', fx);
      for (i = 0; i < 2; i++) {
          if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
              if (i == 0)
                  nums(i);
              if (i == 0 && *out[1])
10                 stars();
              putline(i);
              if (i == 0 && *out[1])
                  fprintf(fx, star);
              if (i == 1)
15                 nums(i);
          }
      }
}

20 /*
   * put out a number line: dumpblock()
   */
   static
   nums(ix)
25     int    ix;      /* index in out[ ] holding seq line */
   {
       char    nline[P_LINE];
       register i, j;
       register char *pn, *px, *py;
30
       for (pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
           *pn = ' ';
       for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
           if (*py == ' ' || *py == '-')
35              *pn = ' ';
           else {
               if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                   j = (i < 0)? -i : i;
                   for (px = pn; j /= 10, px--)
40                      *px = j%10 + '0';
                   if (i < 0)
                       *px = '-';
               }
               else
45                  *pn = ' ';
               i++;
           }
       }
       *pn = '\0';
       nc[ix] = i;
       for (pn = nline; *pn; pn++)
50          (void) putc(*pn, fx);
       (void) putc('\n', fx);
   }
}

55 /*
   * put out a line (name, [num], seq, [num]): dumpblock()
   */
   static
   putline(ix)
60     int    ix;      {

```

...dumpblock

nums

putline

Table 1 (cont')

...putline

```

5      int          i;
      register char *px;

      for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
          (void) putc(*px, fx);
10     for (; i < lmax+P_SPC; i++)
          (void) putc(' ', fx);

      /* these count from 1:
       * ni[] is current element (from 1)
       * nc[] is number at start of current line
       */
15     for (px = out[ix]; *px; px++)
          (void) putc(*px&0x7F, fx);
      (void) putc('\n', fx);
20 }

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
25 static
stars()
{
30     int          i;
      register char *p0, *p1, cx, *px;

      if (!*out[0] || (*out[0] == ' ' && *(p0[0]) == ' ') ||
          !*out[1] || (*out[1] == ' ' && *(p0[1]) == ' '))
          return;
35     px = star;
      for (i = lmax+P_SPC; i; i--)
          *px++ = ' ';

40     for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
          if (isalpha(*p0) && isalpha(*p1)) {

              if (xbm[*p0-'A']&xbm[*p1-'A']) {
                  cx = '*';
                  nm++;
45              }
              else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                  cx = '.';
              else
                  cx = ' ';

50          }
          else
              cx = ' ';

          *px++ = cx;
55      }
      *px++ = '\n';
      *px = '\0';
60 }

```



Table 1 (cont')

```
5  /*
   * strip path or prefix from pn, return len: pr_align0
   */
static
stripname(pn)
    stripname
    char *pn; /* file name (may be path) */
{
10     register char *px, *py;

    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
15             py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
20 }

25

30

35

40

45

50

55

60
```

**Table 1 (cont')**

```

/*
 * cleanup() - cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
5  * g_alloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
10 #include <sys/file.h>

char    *jname = "/tmp/homgXXXXXX";          /* tmp file for jumps */
FILE    *fj;

15 int    cleanup();                          /* cleanup tmp file */
long    lseek();

/*
 * remove any tmp file if we blow
20 */
cleanup(i)
    int    i;
{
    if (fj)
25         (void) unlink(jname);
    exit(i);
}

/*
30 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char    *
35 getseq(file, len)
    char    *file;    /* file name */
    int     *len;      /* seq len */
{
    char    line[1024], *pseq;
    register char    *px, *py;
    int     natgc, tlen;
    FILE    *fp;

    if ((fp = fopen(file, "r")) == 0) {
45         fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }
    tlen = natgc = 0;
    while (fgets(line, 1024, fp)) {
50         if (*line == ';' || *line == '<' || *line == '>')
            continue;
        for (px = line; *px != '\n'; px++)
            if (isupper(*px) || islower(*px))
                tlen++;
55     }
    if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
        fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
        exit(1);
    }
    pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';
60

```

**Table 1 (cont')**

```

py = pseq + 4;
*len = tlen;
rewind(fp);

while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g_calloc(msg, nx, sz)
char *msg;
int nx, sz;
/* program, calling routine */
/* number and size of elements */
{
    char *px, *calloc();

    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g_calloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[ ] or tmp file, set pp[ ], reset dmax: main()
 */
readjmps()
readjmps
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;

    if (fj) {
        (void) fclose(fj);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;

```

Table 1 (cont')

...readjumps

```

5         if (j < 0 && dx[dmax].offset && fj) {
            (void) lseek(fd, dx[dmax].offset, 0);
            (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
            (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
            dx[dmax].ijmp = MAXJMP-1;
        }
10        else
            break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
15    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
20        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;
            /* id = xx - yy + len1 - 1
            */
25            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy++;
            ngapy -= siz;
        /* ignore MAXGAP when doing endgaps */
            siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
30            i1++;
        }
        else if (siz > 0) { /* gap in first seq */
            pp[0].n[i0] = siz;
            pp[0].x[i0] = xx;
35            gapx++;
            ngapx += siz;
        /* ignore MAXGAP when doing endgaps */
            siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
40            i0++;
        }
    }
    else
        break;
}
45
/* reverse the order of jumps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
50     i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
55     i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
60     (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

```

Table 1 (cont')

```

/*
5  * write a filled jmp struct offset of the prev one (if any): nw0
*/
writejumps(ix)
    writejumps
    int    ix;
10  {
    char    *mktemp();

    if (!fj) {
        if (mktemp(jname) < 0) {
15             fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
20             exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
25  }

```

5 PRO XXXXXXXXXXXXXXXX (Length = 15 amino acids)  
Comparison Protein XXXXXYYYYYYY (Length = 12 amino acids)  
% amino acid sequence identity =

**Table 3**

(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%

25    PRO-DNA                                    NNNNNNNNNNNNNNNN                                    (Length = 14 nucleotides)  
       Comparison DNA                        NNNNNNLLLLLLLLLLLL                                    (Length = 16 nucleotides)

      % nucleic acid sequence identity =

### Table 5

66

% nucleic acid sequence identity =

(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) =

5 4 divided by 12 = 33.3%

## II. Compositions and Methods of the Invention

### A. Full-Length PRO Polypeptides

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

As disclosed in the Examples below, various cDNA clones have been disclosed. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

### B. PRO Polypeptide Variants

In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally, the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally

be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

5 PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

10 PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR.

15 Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in

20 reference to amino acid classes, are introduced and the products screened.



Table 6

	Original Residue	Exemplary Substitutions	Preferred Substitutions
5	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
10	Cys (C)	ser	ser
	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
15	Ile (I)	leu; val; met; ala; phe; norleucine	leu
	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
20	Met (M)	leu; phe; ile	leu
	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
25	Trp (W)	tyr; phe	tyr
	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

30 Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- 35 (1) hydrophobic: norleucine, met, ala, val, leu, ile;  
 (2) neutral hydrophilic: cys, ser, thr;  
 (3) acidic: asp, glu;  
 (4) basic: asn, gln, his, lys, arg;  
 (5) residues that influence chain orientation: gly, pro; and  
 40 (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

45 The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487 (1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos.*

Trans. R. Soc. London Ser. A, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

### C. Modifications of PRO

Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propionimide.

Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T.E. Creighton, Proteins: Structure and Molecular Properties, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by,

one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

5 Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 11 September 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

10 Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., Arch. Biochem. Biophys., 259:52 (1987) and by Edge et al., Anal. Biochem., 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., Meth.  
15 Enzymol., 138:350 (1987).

Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

20 The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the PRO. The presence of such epitope-tagged  
25 forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an alpha-tubulin epitope peptide [Skinner et al., J.  
30 Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric

molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

#### D. Preparation of PRO

The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

##### 1. Isolation of DNA Encoding PRO

DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., *supra*; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like <sup>32</sup>P-labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., *supra*.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence

databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., *supra*, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

## 2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., *supra*.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example,  $\text{CaCl}_2$ ,  $\text{CaPO}_4$ , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., *supra*, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*,

*Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kar'*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kar'*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., *J. Bacteriol.*, 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilorum* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., *Proc. Natl. Acad. Sci. USA*, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolytocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., *Biochem. Biophys. Res. Commun.*, 112:284-289 [1983]; Tilburn et al., *Gene*, 26:205-221 [1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479 [1985]). Methylotropic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylotrophs*, 269 (1982).

Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary

(CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

### 3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 $\mu$  plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c)

supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to



300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., Nature, 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

#### 4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

#### 5. Purification of Polypeptide

Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

E. Tissue Distribution

The location of tissues expressing the PRO can be identified by determining mRNA expression in various human tissues. The location of such genes provides information about which tissues are most likely to be affected by the stimulating and inhibiting activities of the PRO polypeptides. The location of a gene in a specific tissue also provides sample tissue for the activity blocking assays discussed below.

As noted before, gene expression in various tissues may be measured by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA (Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 [1980]), dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Gene expression in various tissues, alternatively, may be measured by immunological methods, such as immunohistochemical staining of tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence of a PRO polypeptide or against a synthetic peptide based on the DNA sequences encoding the PRO polypeptide or against an exogenous sequence fused to a DNA encoding a PRO polypeptide and encoding a specific antibody epitope. General techniques for generating antibodies, and special protocols for Northern blotting and *in situ* hybridization are provided below.

F. Antibody Binding Studies

The activity of the PRO polypeptides can be further verified by antibody binding studies, in which the ability of anti-PRO antibodies to inhibit the effect of the PRO polypeptides, respectively, on tissue cells is tested. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies, the preparation of which will be described hereinbelow.

Antibody binding studies may be carried out in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp.147-158 (CRC Press, Inc., 1987).

Competitive binding assays rely on the ability of a labeled standard to compete with the test sample analyte for binding with a limited amount of antibody. The amount of target protein in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies preferably are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected. In a sandwich assay, the test sample analyte is bound by a first antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three-part complex. See, e.g., US Pat No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assay). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

For immunohistochemistry, the tissue sample may be fresh or frozen or may be embedded in paraffin and fixed with a preservative such as formalin, for example.

#### G. Cell-Based Assays

Cell-based assays and animal models for immune related diseases can be used to further understand the relationship between the genes and polypeptides identified herein and the development and pathogenesis of immune related disease.

In a different approach, cells of a cell type known to be involved in a particular immune related disease are transfected with the cDNAs described herein, and the ability of these cDNAs to stimulate or inhibit immune function is analyzed. Suitable cells can be transfected with the desired gene, and monitored for immune function activity. Such transfected cell lines can then be used to test the ability of poly- or monoclonal antibodies or antibody compositions to inhibit or stimulate immune function, for example to modulate monocyte/macrophage proliferation or inflammatory cell infiltration. Cells transfected with the coding sequences of the genes identified herein can further be used to identify drug candidates for the treatment of immune related diseases.

In addition, primary cultures derived from transgenic animals (as described below) can be used in the cell-based assays herein, although stable cell lines are preferred. Techniques to derive continuous cell lines from transgenic animals are well known in the art (see, e.g., Small *et al.*, *Mol. Cell. Biol.* 5: 642-648 [1985]).

The use of an agonist stimulating compound has also been validated experimentally. Activation of 4-1BB by treatment with an agonist anti-4-1BB antibody enhances eradication of tumors. Hellstrom, I. and Hellstrom, K. E., *Crit. Rev. Immunol.* (1998) 18:1. Immunoadjuvant therapy for treatment of tumors, described in more detail below, is another example of the use of the stimulating compounds of the invention.

Alternatively, an immune stimulating or enhancing effect can also be achieved by administration of a PRO which has vascular permeability enhancing properties. Enhanced vascular permeability would be beneficial to disorders which can be attenuated by local infiltration of immune cells (*e.g.*, monocytes/macrophages, eosinophils, PMNs) and inflammation.

5 On the other hand, PRO polypeptides, as well as other compounds of the invention, which are direct inhibitors of monocyte/macrophage proliferation/activation, lymphokine secretion, and/or vascular permeability can be directly used to suppress the immune response. These compounds are useful to reduce the degree of the immune response and to treat immune related diseases characterized by a hyperactive, superoptimal, or autoimmune response. The use of compound which suppress vascular  
10 permeability would be expected to reduce inflammation. Such uses would be beneficial in treating conditions associated with excessive inflammation.

Alternatively, compounds, *e.g.*, antibodies, which bind to stimulating PRO polypeptides and block the stimulating effect of these molecules produce a net inhibitory effect and can be used to suppress the monocyte/macrophage mediated immune response by inhibiting monocyte/macrophage  
15 proliferation/activation and/or lymphokine secretion. Blocking the stimulating effect of the polypeptides suppresses the immune response of the mammal.

#### H. Animal Models

The results of the cell based *in vitro* assays can be further verified using *in vivo* animal models and assays for monocyte/macrophage function. A variety of well known animal models can be used to  
20 further understand the role of the genes identified herein in the development and pathogenesis of immune related disease, and to test the efficacy of candidate therapeutic agents, including antibodies, and other antagonists of the native polypeptides, including small molecule antagonists. The *in vivo* nature of such models makes them predictive of responses in human patients. Animal models of immune related diseases include both non-recombinant and recombinant (transgenic) animals. Non-recombinant animal models  
25 include, for example, rodent, *e.g.*, murine models. Such models can be generated by introducing cells into syngeneic mice using standard techniques, *e.g.*, subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule, *etc.*

Graft-versus-host disease occurs when immunocompetent cells are transplanted into immunosuppressed or tolerant patients. The donor cells recognize and respond to host antigens. The  
30 response can vary from life threatening severe inflammation to mild cases of diarrhea and weight loss. Graft-versus-host disease models provide a means of assessing monocyte/macrophage reactivity against MHC antigens and minor transplant antigens. A suitable procedure is described in detail in Current Protocols in Immunology, above, unit 4.3.

Animal models for delayed type hypersensitivity provides an assay of cell mediated immune  
35 function as well. In chronic Delayed type hypersensitivity (DTH) reactions, monocytes that have differentiated into macrophages lead to the destruction of host tissue which is replaced by fibrous tissue (fibrosis).

Contact hypersensitivity is a simple delayed type hypersensitivity *in vivo* assay of cell mediated immune function. In this procedure, cutaneous exposure to exogenous haptens which gives rise to a delayed type hypersensitivity reaction which is measured and quantitated. Contact sensitivity involves an initial sensitizing phase followed by an elicitation phase. The elicitation phase occurs when the T lymphocytes encounter an antigen to which they have had previous contact. Swelling and inflammation occur, making this an excellent model of human allergic contact dermatitis. At this point, monocytes leave the blood and differentiate into macrophages. A suitable procedure is described in detail in *Current Protocols in Immunology*, Eds. J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach and W. Strober, John Wiley & Sons, Inc., 1994, unit 4.2. See also Grabbe, S. and Schwarz, T., *Immun. Today* 19 (1): 37-44 (1998)

Recombinant (transgenic) animal models can be engineered by introducing the coding portion of the genes identified herein into the genome of animals of interest, using standard techniques for producing transgenic animals. Animals that can serve as a target for transgenic manipulation include, without limitation, mice, rats, rabbits, guinea pigs, sheep, goats, pigs, and non-human primates, *e.g.*, baboons, chimpanzees and monkeys. Techniques known in the art to introduce a transgene into such animals include pronucleic microinjection (Hoppe and Wanger, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (*e.g.*, Van der Putten *et al.*, *Proc. Natl. Acad. Sci. USA* 82, 6148-615 [1985]); gene targeting in embryonic stem cells (Thompson *et al.*, *Cell* 56, 313-321 [1989]); electroporation of embryos (Lo, *Mol. Cel. Biol.* 3, 1803-1814 [1983]); sperm-mediated gene transfer (Lavitrano *et al.*, *Cell* 57, 717-73 [1989]). For review, see, for example, U.S. Patent No. 4,736,866.

For the purpose of the present invention, transgenic animals include those that carry the transgene only in part of their cells ("mosaic animals"). The transgene can be integrated either as a single transgene, or in concatamers, *e.g.*, head-to-head or head-to-tail tandems. Selective introduction of a transgene into a particular cell type is also possible by following, for example, the technique of Lasko *et al.*, *Proc. Natl. Acad. Sci. USA* 89, 6232-636 (1992).

The expression of the transgene in transgenic animals can be monitored by standard techniques. For example, Southern blot analysis or PCR amplification can be used to verify the integration of the transgene. The level of mRNA expression can then be analyzed using techniques such as *in situ* hybridization, Northern blot analysis, PCR, or immunocytochemistry.

The animals may be further examined for signs of immune disease pathology, for example by histological examination to determine infiltration of immune cells into specific tissues. Blocking experiments can also be performed in which the transgenic animals are treated with the compounds of the invention to determine the extent of the monocytes/macrophage proliferation stimulation or inhibition of the compounds. In these experiments, blocking antibodies which bind to the PRO polypeptide, prepared as described above, are administered to the animal and the effect on immune function is determined.

Alternatively, "knock out" animals can be constructed which have a defective or altered gene encoding a polypeptide identified herein, as a result of homologous recombination between the endogenous gene encoding the polypeptide and altered genomic DNA encoding the same polypeptide introduced into

an embryonic cell of the animal. For example, cDNA encoding a particular polypeptide can be used to clone genomic DNA encoding that polypeptide in accordance with established techniques. A portion of the genomic DNA encoding a particular polypeptide can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several  
5 kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see *e.g.*, Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see *e.g.*, Li *et al.*, *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse or  
10 rat) to form aggregation chimeras [see *e.g.*, Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain  
15 the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the polypeptide.

#### I. ImmunoAdjuvant Therapy

In one embodiment, the immunostimulating compounds of the invention can be used in  
20 immunoadjuvant therapy for the treatment of tumors (cancer). It is now well established that monocytes/macrophages recognize human tumor specific antigens. One group of tumor antigens, encoded by the MAGE, BAGE and GAGE families of genes, are silent in all adult normal tissues, but are expressed in significant amounts in tumors, such as melanomas, lung tumors, head and neck tumors, and bladder carcinomas. DeSmet, C. *et al.*, (1996) *Proc. Natl. Acad. Sci. USA*, 93:7149. It has been shown  
25 that stimulation of immune cells induces tumor regression and an antitumor response both *in vitro* and *in vivo*. Melero, I. *et al.*, *Nature Medicine* (1997) 3:682; Kwon, E. D. *et al.*, *Proc. Natl. Acad. Sci. USA* (1997) 94: 8099; Lynch, D. H. *et al.*, *Nature Medicine* (1997) 3:625; Finn, O. J. and Lotze, M. T., *J. Immunol.* (1998) 21:114. The stimulatory compounds of the invention can be administered as adjuvants, alone or together with a growth regulating agent, cytotoxic agent or chemotherapeutic agent, to stimulate  
30 monocyte/macrophage proliferation/activation and an antitumor response to tumor antigens. The growth regulating, cytotoxic, or chemotherapeutic agent may be administered in conventional amounts using known administration regimes. Immunostimulating activity by the compounds of the invention allows reduced amounts of the growth regulating, cytotoxic, or chemotherapeutic agents thereby potentially lowering the toxicity to the patient.

#### J. Screening Assays for Drug Candidates

Screening assays for drug candidates are designed to identify compounds that bind to or complex with the polypeptides encoded by the genes identified herein or a biologically active fragment thereof, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such

screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds, including peptides, preferably soluble peptides, (poly)peptide-immunoglobulin fusions, and, in particular, antibodies including, without limitation, poly-  
5 and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art. All assays are common in that they call for contacting the drug candidate with a polypeptide encoded  
10 by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, *e.g.*, on a microtiter plate, by covalent or non-covalent  
15 attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the polypeptide and drying. Alternatively, an immobilized antibody, *e.g.*, a monoclonal antibody, specific for the polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, *e.g.*, the coated surface containing the anchored component. When  
20 the reaction is complete, the non-reacted components are removed, *e.g.*, by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labelled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a particular protein encoded by a gene identified herein, its interaction with that protein can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields  
30 and co-workers [Fields and Song, *Nature (London)* 340, 245-246 (1989); Chien *et al.*, *Proc. Natl. Acad. Sci. USA* 88, 9578-9582 (1991)] as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA* 89, 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, while the other one functioning as the transcription activation domain. The yeast expression system described in the foregoing publications  
35 (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-*lacZ* reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4

activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for  $\beta$ -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

In order to find compounds that interfere with the interaction of a gene identified herein and other intra- or extracellular components can be tested, a reaction mixture is usually prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a test compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described above. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

#### K. Compositions and Methods for the Treatment of Immune Related Diseases

The compositions useful in the treatment of immune related diseases include, without limitation, proteins, antibodies, small organic molecules, peptides, phosphopeptides, antisense and ribozyme molecules, triple helix molecules, *etc.* that inhibit or stimulate immune function, for example, monocyte proliferation/activation, lymphokine release, or immune cell infiltration.

For example, antisense RNA and RNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation initiation site, *e.g.*, between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, *e.g.*, Rossi, *Current Biology* 4, 469-471 (1994), and PCT publication No. WO 97/33551 (published September 18, 1997).

Nucleic acid molecules in triple helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, *e.g.*, PCT publication No. WO 97/33551, *supra*.

These molecules can be identified by any or any combination of the screening assays discussed above and/or by any other screening techniques well known for those skilled in the art.



## L. Anti-PRO Antibodies

The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

### 1. Polyclonal Antibodies

5 The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or  
10 intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM  
15 adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

### 2. Monoclonal Antibodies

The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature,  
20 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof.  
25 Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells  
30 of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine  
35 ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained,

for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies).

The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Patent No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region

so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

*In vitro* methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

### 3. Human and Humanized Antibodies

The anti-PRO antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*,

Alan R. Liss, p: 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks *et al.*, *Bio/Technology* 10, 779-783 (1992); Lonberg *et al.*, *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild *et al.*, *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

#### 4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, *EMBO J.*, 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh *et al.*, *Methods in Enzymology*, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered

from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan).

Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')<sub>2</sub> bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan *et al.*, Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby *et al.*, J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling *in vitro* to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny *et al.*, J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger *et al.*, Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V<sub>H</sub> and V<sub>L</sub> domains of one fragment are forced to pair with the complementary V<sub>L</sub> and V<sub>H</sub> domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber *et al.*, J. Immunol. 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt *et al.*, *J. Immunol.*, 147:60 (1991).

Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

#### 5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Patent No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared *in vitro* using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

#### 6. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron *et al.*, *J. Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff *et al.*, *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson *et al.*, *Anti-Cancer Drug Design*, 3: 219-230 (1989).

#### 7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain,

nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcun, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta *et al.*, Science, **238**: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (*e.g.*, avidin) that is conjugated to a cytotoxic agent (*e.g.*, a radionucleotide).

#### 8. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein *et al.*, Proc. Natl. Acad. Sci. USA, **82**: 3688 (1985); Hwang *et al.*, Proc. Natl. Acad. Sci. USA, **77**: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin *et al.*, J. Biol. Chem., **257**: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon *et al.*, J. National Cancer Inst., **81**(19): 1484 (1989).

#### 35 M. Pharmaceutical Compositions

The active PRO molecules of the invention (*e.g.*, PRO polypeptides, anti-PRO antibodies, and/or variants of each) as well as other molecules identified by the screening assays disclosed above, can be administered for the treatment of immune related diseases, in the form of pharmaceutical compositions.

Therapeutic formulations of the active PRO molecule, preferably a polypeptide or antibody of the invention, are prepared for storage by mixing the active molecule having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. [1980]), in the form of lyophilized formulations or aqueous solutions.

- 5 Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, 10 gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, 15 Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG).

Compounds identified by the screening assays disclosed herein can be formulated in an analogous manner, using standard techniques well known in the art.

- Lipofections or liposomes can also be used to deliver the PRO molecule into cells. Where 20 antibody fragments are used, the smallest inhibitory fragment which specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable region sequences of an antibody, peptide molecules can be designed which retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology (see, *e.g.*, Marasco *et al.*, *Proc. Natl. Acad. Sci. USA* 90, 7889-7893 [1993]).

- 25 The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise a cytotoxic agent, cytokine or growth inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

- 30 The active PRO molecules may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980). 35

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.



Sustained-release preparations of the PRO molecules may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and  $\gamma$ -ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

#### N. Methods of Treatment

It is contemplated that the polypeptides, antibodies and other active compounds of the present invention may be used to treat various immune related diseases and conditions, such as monocyte/macrophage diseases, including those characterized by infiltration of inflammatory cells into a tissue, stimulation of monocyte/macrophages, inhibition of monocytes/macrophages, increased or decreased vascular permeability or the inhibition thereof.

Exemplary conditions or disorders to be treated with the polypeptides, antibodies and other compounds of the invention, include, but are not limited to systemic lupus erythematosus, rheumatoid arthritis, juvenile chronic arthritis, osteoarthritis, spondyloarthropathies, systemic sclerosis (scleroderma), idiopathic inflammatory myopathies (dermatomyositis, polymyositis), Sjögren's syndrome, systemic vasculitis, sarcoidosis, autoimmune hemolytic anemia (immune pancytopenia, paroxysmal nocturnal hemoglobinuria), autoimmune thrombocytopenia (idiopathic thrombocytopenic purpura, immune-mediated thrombocytopenia), thyroiditis (Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, atrophic thyroiditis), diabetes mellitus, immune-mediated renal disease (glomerulonephritis, tubulointerstitial nephritis), demyelinating diseases of the central and peripheral nervous systems such as multiple sclerosis, idiopathic demyelinating polyneuropathy or Guillain-Barré syndrome, and chronic inflammatory demyelinating polyneuropathy, hepatobiliary diseases such as infectious hepatitis (hepatitis A, B, C, D, E and other non-hepatotropic viruses), autoimmune chronic active hepatitis, primary biliary cirrhosis, granulomatous hepatitis, and sclerosing cholangitis, inflammatory bowel disease (ulcerative colitis: Crohn's disease), gluten-sensitive enteropathy, and Whipple's disease, autoimmune or immune-mediated skin diseases including bullous skin diseases, erythema multiforme and contact dermatitis, psoriasis, allergic diseases such as asthma, allergic rhinitis, atopic dermatitis, food hypersensitivity and

urticaria, immunologic diseases of the lung such as eosinophilic pneumonias, idiopathic pulmonary fibrosis and hypersensitivity pneumonitis, transplantation associated diseases including graft rejection and graft-versus-host-disease.

Rheumatoid arthritis (RA) is a chronic systemic autoimmune inflammatory disease that mainly involves the synovial membrane of multiple joints with resultant injury to the articular cartilage. The pathogenesis is T lymphocyte dependent and is associated with the production of rheumatoid factors, auto-antibodies directed against self IgG, with the resultant formation of immune complexes that attain high levels in joint fluid and blood. These complexes in the joint may induce the marked infiltrate of lymphocytes and monocytes/macrophages into the synovium and subsequent marked synovial changes; the joint space/fluid is infiltrated by similar cells with the addition of numerous neutrophils. Tissues affected are primarily the joints, often in symmetrical pattern. However, extra-articular disease also occurs in two major forms. One form is the development of extra-articular lesions with ongoing progressive joint disease and typical lesions of pulmonary fibrosis, vasculitis, and cutaneous ulcers. The second form of extra-articular disease is the so called Felty's syndrome which occurs late in the RA disease course, sometimes after joint disease has become quiescent, and involves the presence of neutropenia, thrombocytopenia and splenomegaly. This can be accompanied by vasculitis in multiple organs with formations of infarcts, skin ulcers and gangrene. Patients often also develop rheumatoid nodules in the subcutis tissue overlying affected joints; the nodules late stage have necrotic centers surrounded by a mixed inflammatory cell infiltrate. Other manifestations which can occur in RA include: pericarditis, pleuritis, coronary arteritis, interstitial pneumonitis with pulmonary fibrosis, keratoconjunctivitis sicca, and rheumatoid nodules. The number and activation state of macrophages in the inflamed synovium correlates with the significance of RA (Kinne et al., 2000 Arthritis Res. 2: 189-202). As described above, macrophages are not believed to be involved in the early events of RA, but monocytes/macrophages have tissue destructive and tissue remodeling properties which may contribute to both acute and chronic RA.

Juvenile chronic arthritis is a chronic idiopathic inflammatory disease which begins often at less than 16 years of age. Its phenotype has some similarities to RA; some patients which are rheumatoid factor positive are classified as juvenile rheumatoid arthritis. The disease is sub-classified into three major categories: pauciarticular, polyarticular, and systemic. The arthritis can be severe and is typically destructive and leads to joint ankylosis and retarded growth. Other manifestations can include chronic anterior uveitis and systemic amyloidosis.

Spondyloarthropathies are a group of disorders with some common clinical features and the common association with the expression of HLA-B27 gene product. The disorders include: ankylosing spondylitis, Reiter's syndrome (reactive arthritis), arthritis associated with inflammatory bowel disease, spondylitis associated with psoriasis, juvenile onset spondyloarthropathy and undifferentiated spondyloarthropathy. Distinguishing features include sacroileitis with or without spondylitis; inflammatory asymmetric arthritis; association with HLA-B27 (a serologically defined allele of the HLA-B locus of class I MHC); ocular inflammation, and absence of autoantibodies associated with other rheumatoid disease. It was shown that CD163+ macrophages were increased in the synovial lining and

colonic mucosa in Spondyloarthropathy and correlates with the expression of HLA-DR and the production of TNF-alpha (Baeten et al., 2002 J Pathol 196(3):343-350).

Systemic sclerosis (scleroderma) has an unknown etiology. A hallmark of the disease is induration of the skin; likely this is induced by an active inflammatory process. Scleroderma can be localized or systemic; vascular lesions are common and endothelial cell injury in the microvasculature is an early and important event in the development of systemic sclerosis; the vascular injury may be immune mediated. An immunologic basis is implied by the presence of mononuclear cell infiltrates in the cutaneous lesions and the presence of anti-nuclear antibodies in many patients. ICAM-1 is often upregulated on the cell surface of fibroblasts in skin lesions suggesting that T cell interaction with these cells may have a role in the pathogenesis of the disease. As well as T cells, monocytes/macrophages are proposed to play a role in the progression of scleroderma by secreting fibrogenic cytokines (Yamamoto et al., 2001 J Dermatol Sci 26(2): 133-139). Other organs involved include: the gastrointestinal tract: smooth muscle atrophy and fibrosis resulting in abnormal peristalsis/motility; kidney: concentric subendothelial intimal proliferation affecting small arcuate and interlobular arteries with resultant reduced renal cortical blood flow, results in proteinuria, azotemia and hypertension; skeletal muscle: atrophy, interstitial fibrosis; inflammation; lung: interstitial pneumonitis and interstitial fibrosis; and heart: contraction band necrosis, scarring/fibrosis.

Idiopathic inflammatory myopathies including dermatomyositis, polymyositis and others are disorders of chronic muscle inflammation of unknown etiology resulting in muscle weakness. Muscle injury/inflammation is often symmetric and progressive. Autoantibodies are associated with most forms. These myositis-specific autoantibodies are directed against and inhibit the function of components, proteins and RNA's, involved in protein synthesis.

Sjögren's syndrome is due to immune-mediated inflammation and subsequent functional destruction of the tear glands and salivary glands. The disease can be associated with or accompanied by inflammatory connective tissue diseases. The disease is associated with autoantibody production against Ro and La antigens, both of which are small RNA-protein complexes. Lesions result in keratoconjunctivitis sicca, xerostomia, with other manifestations or associations including biliary cirrhosis, peripheral or sensory neuropathy, and palpable purpura.

Systemic vasculitis are diseases in which the primary lesion is inflammation and subsequent damage to blood vessels which results in ischemia/necrosis/degeneration to tissues supplied by the affected vessels and eventual end-organ dysfunction in some cases. Vasculitis can also occur as a secondary lesion or sequelae to other immune-inflammatory mediated diseases such as rheumatoid arthritis, systemic sclerosis, etc., particularly in diseases also associated with the formation of immune complexes. Diseases in the primary systemic vasculitis group include: systemic necrotizing vasculitis: polyarteritis nodosa, allergic angiitis and granulomatosis, polyangiitis; Wegener's granulomatosis; lymphomatoid granulomatosis; and giant cell arteritis. Miscellaneous vasculitides include: mucocutaneous lymph node syndrome (MLNS or Kawasaki's disease), isolated CNS vasculitis, Behet's disease, thromboangiitis obliterans (Buerger's disease) and cutaneous necrotizing venulitis. The pathogenic mechanism of most of

the types of vasculitis listed is believed to be primarily due to the deposition of immunoglobulin complexes in the vessel wall and subsequent induction of an inflammatory response either via ADCC, complement activation, or both.

5 Sarcoidosis is a condition of unknown etiology which is characterized by the presence of epithelioid granulomas in nearly any tissue in the body; involvement of the lung is most common. The pathogenesis involves the persistence of activated macrophages and lymphoid cells at sites of the disease with subsequent chronic sequelae resultant from the release of locally and systemically active products released by these cell types.

10 Autoimmune hemolytic anemia including autoimmune hemolytic anemia, immune pancytopenia, and paroxysmal nocturnal hemoglobinuria is a result of production of antibodies that react with antigens expressed on the surface of red blood cells (and in some cases other blood cells including platelets as well) and is a reflection of the removal of those antibody coated cells via complement mediated lysis and/or ADCC/Fc-receptor-mediated mechanisms.

15 Thyroiditis including Grave's disease, Hashimoto's thyroiditis, juvenile lymphocytic thyroiditis, and atrophic thyroiditis, are the result of an autoimmune response against thyroid antigens with production of antibodies that react with proteins present in and often specific for the thyroid gland. Experimental models exist including spontaneous models: rats (BUF and BB rats) and chickens (obese chicken strain); inducible models: immunization of animals with either thyroglobulin, thyroid microsomal antigen (thyroid peroxidase).

20 Inflammatory and Fibrotic Lung Disease, including Eosinophilic Pneumonias; Idiopathic Pulmonary Fibrosis, and Hypersensitivity Pneumonitis may involve a dysregulated immune-inflammatory response. Inhibition of that response would be of therapeutic benefit.

Psoriasis is a T lymphocyte-mediated inflammatory disease. Lesions contain infiltrates of T lymphocytes, macrophages and antigen processing cells, and some neutrophils.

25 Other diseases in which intervention of the immune and/or inflammatory response have benefit are infectious disease including but not limited to viral infection (including but not limited to AIDS, hepatitis A, B, C, D, E and herpes) bacterial infection, fungal infections, and protozoal and parasitic infections. Molecules (or derivatives/agonists) which stimulate the immune reaction can be utilized therapeutically to enhance the immune response to infectious agents), diseases of immunodeficiency  
30 (molecules/derivatives/agonists) which stimulate the immune reaction can be utilized therapeutically to enhance the immune response for conditions of inherited, acquired, infectious induced (as in HIV infection), or iatrogenic (*i.e.*, as from chemotherapy) immunodeficiency, and neoplasia.

It has been demonstrated that some human cancer patients develop an antibody and/or monocyte/macrophage response to antigens on neoplastic cells. It has also been shown in animal models  
35 of neoplasia that enhancement of the immune response can result in rejection or regression of that particular neoplasm. Molecules that enhance the monocyte/macrophage response have utility *in vivo* in enhancing the immune response against neoplasia. Molecules which enhance the monocyte/macrophage proliferative response (or small molecule agonists or antibodies that affected the same receptor in an

agonistic fashion) can be used therapeutically to treat cancer. Molecules that inhibit the monocyte/macrophage response also function *in vivo* during neoplasia to suppress the immune response to a neoplasm; such molecules can either be expressed by the neoplastic cells themselves or their expression can be induced by the neoplasm in other cells. Antagonism of such inhibitory molecules (either with antibody, small molecule antagonists or other means) enhances immune-mediated tumor rejection.

Additionally, inhibition of molecules with proinflammatory properties may have therapeutic benefit in reperfusion injury; stroke; myocardial infarction; atherosclerosis; acute lung injury; hemorrhagic shock; burn; sepsis/septic shock; acute tubular necrosis; endometriosis; degenerative joint disease and pancreatitis.

The compounds of the present invention, *e.g.*, polypeptides or antibodies, are administered to a mammal, preferably a human, in accord with known methods, such as intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation (intranasal, intrapulmonary) routes. Intravenous or inhaled administration of polypeptides and antibodies is preferred.

In immunoadjuvant therapy, other therapeutic regimens, such administration of an anti-cancer agent, may be combined with the administration of the proteins, antibodies or compounds of the instant invention. For example, the patient to be treated with a the immunoadjuvant of the invention may also receive an anti-cancer agent (chemotherapeutic agent) or radiation therapy. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in *Chemotherapy Service* Ed., M.C. Perry, Williams & Wilkins, Baltimore, MD (1992). The chemotherapeutic agent may precede, or follow administration of the immunoadjuvant or may be given simultaneously therewith. Additionally, an anti-estrogen compound such as tamoxifen or an anti-progesterone such as onapristone (see, EP 616812) may be given in dosages known for such molecules.

It may be desirable to also administer antibodies against other immune disease associated or tumor associated antigens, such as antibodies which bind to CD20, CD11a, CD18, ErbB2, EGFR, ErbB3, ErbB4, or vascular endothelial factor (VEGF). Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be coadministered to the patient. Sometimes, it may be beneficial to also administer one or more cytokines to the patient. In one embodiment, the PRO polypeptides are coadministered with a growth inhibitory agent. For example, the growth inhibitory agent may be administered first, followed by a PRO polypeptide. However, simultaneous administration or administration first is also contemplated. Suitable dosages for the growth inhibitory agent are those presently used and may be lowered due to the combined action (synergy) of the growth inhibitory agent and the PRO polypeptide.

For the treatment or reduction in the severity of immune related disease, the appropriate dosage of an a compound of the invention will depend on the type of disease to be treated, as defined above, the

severity and course of the disease, whether the agent is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the compound, and the discretion of the attending physician. The compound is suitably administered to the patient at one time or over a series of treatments.

- 5 For example, depending on the type and severity of the disease, about 1  $\mu\text{g/kg}$  to 15  $\text{mg/kg}$  (*e.g.*, 0.1-20  $\text{mg/kg}$ ) of polypeptide or antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about 1  $\mu\text{g/kg}$  to 100  $\text{mg/kg}$  or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is  
10 sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

O. Articles of Manufacture

- In another embodiment of the invention, an article of manufacture containing materials (*e.g.*, comprising a PRO molecule) useful for the diagnosis or treatment of the disorders described above is  
15 provided. The article of manufacture comprises a container and an instruction. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for diagnosing or treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The  
20 active agent in the composition is usually a polypeptide or an antibody of the invention. An instruction or label on, or associated with, the container indicates that the composition is used for diagnosing or treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint,  
25 including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

P. Diagnosis and Prognosis of Immune Related Disease

- Cell surface proteins, such as proteins which are overexpressed in certain immune related diseases, are excellent targets for drug candidates or disease treatment. The same proteins along with secreted proteins encoded by the genes amplified in immune related disease states find additional use in the  
30 diagnosis and prognosis of these diseases. For example, antibodies directed against the protein products of genes amplified in multiple sclerosis, rheumatoid arthritis, or another immune related disease, can be used as diagnostics or prognostics.

- For example, antibodies, including antibody fragments, can be used to qualitatively or quantitatively detect the expression of proteins encoded by amplified or overexpressed genes ("marker  
35 gene products"). The antibody preferably is equipped with a detectable, *e.g.*, fluorescent label, and binding can be monitored by light microscopy, flow cytometry, fluorimetry, or other techniques known in the art. These techniques are particularly suitable, if the overexpressed gene encodes a cell surface protein. Such binding assays are performed essentially as described above.

*In situ* detection of antibody binding to the marker gene products can be performed, for example, by immunofluorescence or immunoelectron microscopy. For this purpose, a histological specimen is removed from the patient, and a labeled antibody is applied to it, preferably by overlaying the antibody on a biological sample. This procedure also allows for determining the distribution of the marker gene product in the tissue examined. It will be apparent for those skilled in the art that a wide variety of histological methods are readily available for *in situ* detection.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

#### EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA.

##### EXAMPLE 1: Microarray analysis of monocyte/macrophages.

Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in diseased tissues as compared to their normal counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. For example, a selection of genes known to be expressed in certain disease states may be arrayed on a solid support. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (in this instance, differentiated macrophages) sample is greater than hybridization signal of a probe from a control (in this instance, non-differentiated monocytes) sample, the gene or genes expressed in the test tissue are identified. The implication of this result is that an overexpressed protein in a test tissue is useful not only as a diagnostic marker for the presence of the disease condition, but also as a therapeutic target for treatment of the disease condition.

The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In one example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in PCT Patent Application Serial No. PCT/US01/10482, filed on March 30, 2001 and which is herein incorporated by reference.

In this experiment, CD14+ monocytes are selected by positive selection according to Miltenyi MACS™ protocol. Lymphocytes in 100 ml heparinized blood are separated using Ficoll Paque™. Cells are washed twice in PBS/0.5% BSA/2 mM EDTA. In final wash, all gradients are pooled and volume is brought to approximately 10 ml. The cells are centrifuged, the supernatant is removed and the cell pellet

is resuspended in buffer in a total volume of  $10^7$  cells per 80  $\mu$ l buffer. Add 20  $\mu$ l CD14 microbeads per  $10^7$  total cells, mix and incubate 15 minutes at 6-12 C. Wash the cells by adding 20x labeling volume of buffer, spin pellet and resuspend in 500  $\mu$ l buffer per  $10^8$  cells. Separate cells with MACS™ depletion column type D and check purity of cells by labeling with anti-CD45 and anti-CD14 antibodies (cell purity at this point is >95%). Lyse cells in RNA lysis buffer to obtain a timepoint of Day 0 monocytes, then plate remaining cells in 6 well plates in macrophage differentiation medium: DMEM 4.5  $\mu$ g/ml glucose, Pen-Strep, L-glutamine, 20% FBS and 10% Human AB serum (Gemini, Cat # 100-512). Seed cells at  $1.5 \times 10^6$  per well (6 well Costar cell culture plates) and grow at 37 C, 7% CO<sub>2</sub>. After 24 hours in culture, the cells were harvested and lysed in RNA lysis buffer to obtain mRNA for the Day 1 timepoint. The remaining cells were kept in culture and until Day 7. After 7 days in culture, the cells were lysed in RNA lysis buffer to obtain Day 7 timepoint at which time the cells displayed gross macrophage morphology.

The mRNA was isolated by Qiagen miniprep and analysis run on Affimax™ (Affymetrix Inc. Santa Clara, CA) microarray chips and proprietary Genentech microarrays. The cells harvested at Day 0 timepoint, the Day 1 timepoint, and the Day 7 timepoint were subjected to the same analysis. Genes were compared whose expression was upregulated at Day 7 as compared to Day 0 and Day 1.

Below are the results of these experiments, demonstrating that various PRO polypeptides of the present invention are differentially expressed in differentiated macrophages at Day 7 as compared to non-differentiated monocytes at Day 0 and at Day 1. As described above, these data demonstrate that the PRO polypeptides of the present invention are useful not only as diagnostic markers for the presence of one or more immune disorders, but also serve as therapeutic targets for the treatment of those immune disorders. Specifically, the cDNAs shown Figures 592, Figure 708, Figure 724, Figure 888, Figure 1095, Figure 1109, Figure 1456 and Figure 2331 are significantly overexpressed in differentiated macrophages as compared to non-differentiated monocytes at Day 0 and Day 1.

The Figures 1-2517 show the nucleic acids of the invention and their encoded PRO polypeptides that are differentially expressed in differentiated macrophages at Day 7 as compared to non-differentiated monocytes at Day 0 and at Day 1.

#### EXAMPLE 2: Use of PRO as a hybridization probe

The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.



DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

EXAMPLE 3: Expression of PRO in *E. coli*

5 This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable  
10 vector is pBR322 (derived from *E. coli*; see Bolivar et al., Gene, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO  
15 coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

20 Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the  
25 solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector,  
30 and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an O.D.<sub>600</sub> of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.71 g sodium  
35 citrate•2H<sub>2</sub>O, 1.07 g KCl, 5 .36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO<sub>4</sub>) and grown for approximately

20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

*E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### EXAMPLE 4: Expression of PRO in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO.

5 In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10  $\mu$ g pRK5-PRO DNA is mixed with about 1  $\mu$ g DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500  $\mu$ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M  $\text{CaCl}_2$ . To this mixture is added,  
10 dropwise, 500  $\mu$ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM  $\text{NaPO}_4$ , and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

15 Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200  $\mu\text{Ci/ml}$   $^{35}\text{S}$ -cysteine and 200  $\mu\text{Ci/ml}$   $^{35}\text{S}$ -methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may  
20 undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700  $\mu$ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate  
25 is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5  $\mu\text{g/ml}$  bovine insulin and 0.1  $\mu\text{g/ml}$  bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column  
30 chromatography.

In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as  $\text{CaPO}_4$  or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as  $^{35}\text{S}$ -methionine. After determining the presence of PRO polypeptide, the culture  
35 medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 promoter/enhancer containing vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 promoter/enhancer containing vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by  $\text{Ni}^{2+}$ -chelate affinity chromatography.

PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., Nucl. Acids Res. 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect<sup>®</sup> (Quiagen), Dospers<sup>®</sup> or Eugene<sup>®</sup> (Boehringer Mannheim). The cells are grown as described in Lucas et al., supra. Approximately  $3 \times 10^7$  cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mL of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2  $\mu\text{m}$  filtered PS20 with 5% 0.2  $\mu\text{m}$  diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37°C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with  $3 \times 10^5$  cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 may actually be used. A 3L production spinner is seeded at  $1.2 \times 10^6$  cells/mL. On day 0, pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 mL of 500 g/L glucose and 0.6

mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical-Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22  $\mu$ m filter. The filtrate was either stored at 4°C or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275  $\mu$ l of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### EXAMPLE 5: Expression of PRO in Yeast

The following method describes recombinant expression of PRO in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### EXAMPLE 6: Expression of PRO in Baculovirus-Infected Insect Cells

The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (PharMingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

Expressed poly-his tagged PRO can then be purified, for example, by Ni<sup>2+</sup>-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., Nature, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl<sub>2</sub>; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 µm filter. A Ni<sup>2+</sup>-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A<sub>280</sub> with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A<sub>280</sub> baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni<sup>2+</sup>-NTA-conjugated to alkaline phosphatase

(Qiagen). Fractions containing the eluted His<sub>10</sub>-tagged PRO are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

5 Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

#### EXAMPLE 7: Preparation of Antibodies that Bind PRO

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

10 Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms.

15 Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-  
20 PRO antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The  
25 fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in  
30 the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion  
35 chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 8: Purification of PRO Polypeptides Using Specific Antibodies

Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (*e.g.*, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (*e.g.*, a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

EXAMPLE 9: Drug Screening

This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested.

Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an



agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

#### EXAMPLE 10: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

In one approach, the three-dimensional structure of the PRO polypeptide, or of a PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon

which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**